

Investigation of the Effects of Carbonation and Orange Juice on the Physical, Chemical and Microbial Characteristics of Pasteurized Carrot Juice

Hatami, Z.¹, Hamidi-Esfahani, Z.^{2*}, Abbasi, S.³

1-M.Sc. Graduate, Department of Food Science and Technology, Faculty of Agriculture, Tarbiat Modares University

2-Associate Prof., Department of Food Science and Technology, Faculty of Agriculture, Tarbiat Modares University

3- Assistant Prof., Department of Food Science and Technology, Faculty of Agriculture, Tarbiat Modares University

Abstract

The effect of CO₂ (0.5 l min⁻¹ for 15 min) and orange juice (30% v/v carrot juice) on physico-chemical characteristics as well as microbial flora of carrot juice within the three months of storage was investigated. Both orange juice and CO₂ caused a decrease in pH value of carrot juice but pH reduction was higher when orange juice (30 percent) was added. Samples of carrot-orange juice had higher value of acidity and lower pH. Blending orange juice with carrot juice caused higher percentage of precipitation, decrease in a* and b* values and increase in L* value. The results of this experiment showed that sugar decreases the percentage of precipitation in pasteurized carrot juice. No microbial growth was detected in either of the treatments.

Keywords: Carrot-orange juice, Carbonation, Physico-chemical, Storage time, Microbial, Sugar content

1-Introduction

Fruit and vegetable juices are recommended for a healthy diet and various health benefits [1]. Fruit juices are rich sources of vitamin C, E, phenolic compounds, carotenoids and can prevent type of certain diseases (cancer and cardiovascular) [2-6].

There are different treatments which makes juices more acceptable for consumers. Beverages containing carbon dioxide are very popular products nowadays [7-8] and carbonated carrot juice can be appreciable as one of these products. Carbon dioxide enhances the taste of beverages and their appearance [9]. In addition, CO₂ can cause a decrease in pH and at high pressures

has an inhibitory effects on microbial growth [10] but high levels of gas spoil the flavour of fruit and vegetable juices [11]. Blending with orange juice can be considered as another treatment of carrot juice that in addition to induce the special sensory and nutritive characteristics, it provides a good source of functional compounds, vitamins and antioxidants that contribute to the health of the consumers [12]. The growth of some bacteria, mold and yeast flora in mixed orange and carrot juice [13-15] and the effect of PEF (pulsed electric field) and heat pasteurization on some physical-chemical characteristics of carrot-orange juice [16] have been

*Corresponding author E-mail address: hamidy_z@modares.ac.ir

investigated up to now.

The objective of this study was to evaluate the influence of CO₂, orange juice and sugar addition on some chemical, physical, sensorial and microbial characteristics of pasteurized carrot juice.

2-Materials and Methods

2-1-Treatment of Carrot Juice

Fresh carrots (*Daucus carrota* L. cv. Alpha) of good quality were purchased from a local market and stored at 5 °C. Juices were extracted and prepared as follows:

After peeling and washing with water (manual), the carrots were grinded with grinding mill (Kenwood FP/690, England). As acidification before juice extraction can prevent clarification of carrot juice, citric acid (0.5 g kg⁻¹) (BBCA company, Arhui, China) was added and then the mash was heated (70 °C, 10 min) in water bath (Memmert, WB14, Germany) for blanching. To increase the yield of juice extraction, the enzymes of pectinase (Rohapect PTE) and cellulase (Rohament CL) prepared from AB-Enzyme Company (Darmstadt, Germany) were added and mash incubation was performed at 40 °C for 45 minutes in incubator (FOC 225I, Europe). After manual juice extraction, clarification was done by centrifuge at 3000 rpm for 10 minutes (Kubota 6900, Japan). Pomace was separated and the juice was pasteurized (95 °C, 1 min) in rotary (Heidolph 4001, Germany) [17-18].

Concentrated orange juice with °Brix of 70 was purchased from Ramsar Gardeners Factory (Ramsar, Iran) and reconstituted with water to obtain a regular orange juice. To prepare carrot-orange juice blend, carrot (70%), orange (30%) and sugar (4%) were mixed.

For treating pasteurized carrot juice with CO₂, a glass instrument with a cold water jacket was designed (Fig. 1).

Since CO₂ dissolves better at low temperatures [7] the jacket was designed. CO₂ was bubbled through carrot juice from a gas cylinder via a plastic tube at a rate of approximately 0.5 l min⁻¹ for 15 minutes.

The juice was then bottled in sterilized glass bottles with screw caps and they were stored at 4 °C for three months.




Fig. 1. Carbonation unit: (1) CO₂ cylinder; (2) carrot juice; (3) water inlet; (4) water outlet; (5) gas outlet; (6) glass tank.

A total of four juice treatments were produced to compare with carrot juice alone (C) as control. Treatments were: the pasteurized carrot juice with 4% sugar (CS), the pasteurized carrot juice treated with CO₂ (CC), the pasteurized carrot juice with 4% sugar and the samples treated with CO₂ (CSC) and the mixed carrot orange juice including 4% sugar (COS). All the samples were stored at 4 °C and their physico-chemical and microbiological characteristics were investigated at the first of storage and during three months of storage with one week intervals.

2-2-Physico-chemical Analysis

pH measurements were performed using a Metrohm model 744 pH meter (Switzerland). Titrable acidity was determined potentiometrically using 0.1 N NaOH and phenolphthalein (indicator). The results were expressed as g 100 ml⁻¹ on the basis of citric acid. Total soluble solids (TSS) was measured by the refractive index in terms of °Brix with an Atago N1 refractometer (Japan) [19]. Percentage of

precipitation was determined after calculating the height of sediment that accumulated at the bottom of the test tubes. Color of the juices was measured by Hunter Lab model Colorflex 40/0 (USA) in mode of Daylight color. Letters L^* , a^* , and b^* represent luminosity or lightness, redness and yellowness, respectively [20].

Each sample was assessed at the first of storage and for three months of storage with one week intervals except color which was measured with one month intervals.

2-3-Microbial Analysis

To evaluate the microbiological efficiency of the treatments, one ml of each sample without dilution was cultured (1ml/plate) to determine the number of colony forming units (CFU/ml) using pour plate technique. Media of plate count agar (Merck) for aerobic mesophyl bacteria, orange serum agar (Merck) for acidophyl bacteria and yeast extract glucose chloramphenicol agar (Merck) for yeasts and moulds were used [21].

2-4-Sensory Analysis

25 untrained volunteers were selected for sensory analysis. The samples of carbonated pasteurized carrot juice with 4% sugar (CSC), carrot orange juice blend with 4% sugar (COS) along with carrot juice alone (C) as control were coded and after ranking the samples by assessors, preferred sample was determined [22-23].

2-5-Statistical Analysis

Experimental data were analyzed by analysis of variance (ANOVA) with significance defined at $P < 0.05$, and significance differences between the means were determined by Duncan's multiple range test. SPSS software version 12.0 was performed for the statistical analysis. All the reported values are the means of triplicate determinations.

3-Results and Discussion

The effect of the treatments on chemical (pH, acidity, °Brix, percentage of precipitation, color) and microbial flora were as follows:

3-1-Effect on pH

Results showed that sample of blended carrot and orange juice (COS) had the lowest pH value because of lower pH of orange juice and the samples treated with CO_2 (with and without sugar) were in the same group without any significant difference (Fig. 2). These treatments with significant difference had lower pH value than the control sample and carrot juice with 4% sugar. Alkint *et al.* (2004), also reported that adding CO_2 bubbles decreases the pH of carrot juice [24].

There was no pH variation during storage period which is in good agreement with the results reported for blended orange and carrot juice (80% orange and 20% carrot) [16], pasteurized carrot juice [25], and orange juice treated with Pulsed Electric Fields (PEF) [26].




Fig. 2. Effects of different treatments on pH. Different letters show significant difference between the treatments ($p < 0.05$) by duncan's multiple range test.

3-2-Effect on Titrable Acidity

With reference to our findings (Fig. 3) all the samples were located in two groups. Among the samples, the highest acidity allocated to treatment of blended carrot and orange juice (COS) which is because of higher acidity nature of orange. In another word, this sample had the lowest pH and the highest acidity.

Although the titrable acidity of CSC and CC treatments were slightly higher than the

CS and C samples but there was no significant difference between them, statistically. According to Woodroof and Philips (1981) findings, carbonation increased the sour flavor and titrable acidity of treatments including grape juice and blended grape and apple juice [27]. The acidity of samples did not vary ($p < 0.05$) during storage but Rivas *et al.* 2006 recorded that during storage, the sample of blended orange and carrot juice treated with PEF showed an increase in total acidity after 7.5 weeks of storage at 12 °C [16].




Fig. 3. Effects of different treatments on titrable acidity. Different letters show significant difference between the treatments ($p < 0.05$) by duncan's multiple range test.

3-3-Effect on Total Soluble Solids

With reference to information of variance analysis (not shown) and Fig. 4 all treatments had significant difference. It can be mentioned that because of higher content of soluble solids, especially sugars of carrot juice, treatments which contain carrot and sugar showed the greatest °brix. As orange juice has less soluble solids than carrot juice, with participation of orange in a mixed juice, the °brix decreased.

In fruit and vegetable juices, fermentation of sugars by microorganisms can cause spoilage. Therefore, °brix will change. Due to absence of microbial spoilage in our treatments, there were no significant changes in °brix. These findings are similar

to the ones reported by Rivas *et al.* 2006 and Yoem *et al.* 2000 [16, 26].




Fig. 4. Effects of different treatments on °brix. Different letters show significant difference between the treatments ($p < 0.05$) by duncan's multiple range test.

3-4-Effect on Percentage of Precipitation

As it can be seen (Fig. 5), treatments of carrot juice which contained 4% sugar and carbonated carrot juice with 4% sugar had lower percentage of precipitation. This behavior seems to reflect the thickening effect of sugar in solutions such as juices [28]. But the sample of COS had the greatest percentage of precipitation. As orange juice has a high acidity, after mixing it with carrot juice, the proteins were coagulated and their tendency to precipitation increased.




Fig. 5. Effects of different treatments on percentage of precipitation. Different letters show significant difference between the treatments ($p < 0.05$) by duncan's multiple range test.

3-5-Effect on Color

To evaluate of color three parameters (L^* , a^* and b^*) that represent lightness, redness and yellowness, respectively were measured (Fig. 6). The highest L^* and lowest a^* was related to COS treatment. Citrus have great lightness [29], as a result the L^* of mixture of orange-carrot juice was higher than other treatments. Since β -caroten is the major carotenoid (60–80%) of carrot pigments which is responsible for orange-red color, with decreasing of carrot in mixed carrot orange juice, the a^* value decreased [30]. With regard to the effect of storage at 4 °C for 3 months on color parameters, no significant variation was found. It needs to be mentioned that also Rivas *et al.* 2006 found the same result for L^* of the PEF-treated samples and opposite result for L^* of pasteurized blended orange and carrot juice in which it diminished (at 12°C) [16].






Fig. 6. Effects of different treatments on color parameters: L^* (a), a^* (b) and b^* (c). Different letters show significant difference between the treatments ($p < 0.05$) by duncan's multiple range test

3-6-Sensory Analysis

As it shown in Fig. 7, the preferred sample devoted to carbonated carrot juice with 4% sugar. It may be worth to say that this results can be different in different region because of people appetite.




Fig. 7. Effect of different treatment of pasteurized carrot orange blend including 4% sugar (COS), pasteurized carrot juice (C) and carbonated pasteurized carrot juice including 4% sugar (CSC) on pleasuring factor.

3-7-Microbial Analysis

During the time of this experiment no microbial growth including aerophil bacteria, acidophil bacteria, yeast and mold was observed.

4-Conclusions

On the basis of the results presented it may be concluded that both orange juice and CO₂ cause decrease in pH value and increase in acidity of carrot juice. The highest precipitation related to mixture of carrot and orange juice and in contrast samples treated with sugar and sugar along with CO₂ had the lowest precipitation. It can be mentioned that as the in prolonged shelf life, the precipitation also will increase. The results of this experiment showed that addition of sugar decreases the precipitation percentage of pasteurized carrot juice and addition of orange juice increases the L^* value and decreases the a^* and b^* values. Most probably it is due to the content of pigments in orange juice. During the time of this experiment no microbial growth was observed.

5-Acknowledgement

The authors would like to thank the Tarbiat Modares University for financial support.

6-References

- [1] Song, H.P., Kim, D.H., Jo, Ch., Lee, Ch.H., Kim, K.S. and Byun, M.W. 2006. Effect of gamma irradiation on the microbiological quality and antioxidant activity of fresh vegetable juice. *Food Microbiol.* 23, 372–378.
- [2] Burns, J., Frase, P.D. and Bramley, P.M. 2003. Identification and quantification of carotenoids, tocopherols and chlorophylls in commonly consumed fruits and vegetables. *Phytochem.* 62, 939–947.
- [3] Gardner, P.T., White, T.A.C., Mcphail, D.B. and Duthie, G.G. 2000. The relative contributions of vitamin C, carotenoids and phenolics to the antioxidant potential of fruit juices. *Food Chem.* 68, 471–474.
- [4] John, J.H., Ziebland, S., Yudkin, P., Roe, L.S. and Neil, H.A.W. 2002. Effects of fruit and vegetable consumption on plasma antioxidant concentrations and blood pressure: A randomised controlled trial. *Lancet*, 359, 1969–1974.
- [5] Rubatzky, V.E. and Yamaguchi, M. 1997. *World Vegetables*, 2nd Ed., Capman & Hall, New York.
- [6] Gross, J. 1991. *Pigments in Vegetables*, Van Nostrand Reinhold, New York.
- [7] Descoins, Ch., Mathlouthi, M., Moual, M.L. and Hennequin, J. 2006. Carbonation monitoring of beverage in a laboratory scale unit with on-line measurement of dissolved CO₂. *Food Chem.* 95, 541–553.
- [8] Otake, S. 2001. Sweetness intensity in low-carbonated beverages. *Biomol. Eng.* 17, 151–156.
- [9] Kupper, L. Philip and Marsha, W., 1986. United State Patent 4,612,205.
- [10] Molin, G. 2000. Modified atmosphere. In *The microbiological safety and quality of food*, (B.M. Lund, T.C. BairdParker and G.W. Gould, eds). pp. 214–234. Aspen Publishers, Inc, Gaithersburg.
- [11] Barker, G., Jefferson, B. and Judd, S.J. 2002. Domestic carbonation process optimisation. *J. Food Eng.* 52, 405–412.
- [12] Torregrosa, F., Esteve, M.J., Frígola, A. and Cortés, C. 2006. Ascorbic acid stability during refrigerated storage of orange-carrot juice treated by high pulsed electric field and comparison with pasteurized juice. *J. Food Eng.* 73, 339–345.
- [13] Rodrigo, D., Barbosa-Cánovas, G.V., Martínez, A. and Rodrigo, M. 2003. Weibull distribution function based on an empirical mathematical model for inactivation of *Escherichia coli* by pulsed electric fields. *J. Food Protect.* 66, 1007–1012.
- [14] Rodrigo, D., Barbosa-Cánovas, G.V., Martínez, A. and Rodrigo, M. 2003b. Pectin methyl esterase and natural microbial flora of fresh mixed orange and carrot juice treated with pulsed electric fields. *J. Food Protect.* 66, 2336–2342.
- [15] Rodrigo, D., Martínez, A., Harte, F., Barbosa-Cánovas, G.V. and Rodrigo, M. 2001. Study of inactivation of *Lactobacillus plantarum* in orange-carrot juice by means of pulsed electric fields: Comparison of inactivation kinetics models. *J. Food Protect.* 64, 259–263.

- [16] Rivas, A., Rodrigo, D., Martinez, A., Barbosa-Cánovas, G. V. and Rodrigo, M. 2006. Effect of PEF and heat pasteurization on the physical-chemical characteristics of blended orange and carrot juice. *Food Sci. Technol.* 39, 1163–1170.
- [17] Mokhtari, Z., Hamidi, Z., and Azizi, M.H. 2005. Effect of various processing methods on the cloud stability of carrot juice. *Iran. J. food Sci. Technol.* 1, 27– 33.
- [18] Hamidi-Esfahani, Z., Mokhtary, Z., and Azizi, M.H. 2006. Study on optimization for carrot juice production. *Proceedings of the Young Researchers Conference on Applied Science*, Shah Alam, Malaysia.
- [19] Anonymous. 1988. Fruit juices-test methods. Institute of Standards and Industrial Research of Iran. No, 2685.
- [20] Chen, B.H., Peng, H.Y. and Chen, H.E. 1995. Changes of color and vitamin A content during processing of carrot juice. *J. Agr. Food Chem.* 43, 1972–1978.
- [21] Anonymous. 1994. Microbiological characteristics and estimation of hygiening conditions of fruit products prevented only by physical methods. Institute of Standards and Industrial Research of Iran. No, 3414.
- [22] Anonymous. 1993. Sensorial test methods-knowledge general guide of sensorial test methods. Institute of Standards and Industrial Research of Iran. No, 3442.
- [23] Anonymous. 1993. Sensorial test methods-evaluation of nutritious products with comparative methods. Institute of Standards and Industrial Research of Iran. No, 3443.
- [24] Alkint, C., Wadsö, L. and Sjöholm, I. 2004. Effects of modified atmosphere on shelf-life of carrot juice. *Food Control.* 15, 131–137.
- [25] Saldana, G., Stephens, T.S. and Lime, B.J. 1976. Carrot beverages. *J. Food Sci.* 41, 1243-1244.
- [26] Yoem, H.W., Streaker, C.B., Zhang, Q.H., and Min, D.B. 2000. Effects of pulsed electric fields on the quality of orange juice and comparison with heat pasteurization. *J. Agr. Food Chem.* 48, 4597–4605.
- [27] Woodroof, J.G., and Phillips, G.F. 1981. *Beverages: Carbonated and uncarbonated*, AVI Publishing Company Incorporated, USA.
- [28] Shachman, M. 2005. *The Soft Drinks Companion: A Technical Handbook for the Beverage Industry*, CRC Press, USA.
- [29] Kimball, D. 1999. *Citrus Processing*, 2nd Ed., Chapman & Hall, California.
- [30] Sun, M., and Temelli, F. 2006. Supercritical carbon dioxide extraction of carotenoids from carrot using canola oil as a continuous co-solvent. *J. Supercrit. Fluids.* 37, 397-408.

بررسی تأثیرات کربناته کردن و آب پرتقال بر خصوصیات فیزیکی، شیمیایی و میکروبی آب هویج پاستوریزه

زینب حاتمی^۱، زهره حمیدی اصفهانی^{۲*}، سلیمان عباسی^۳

۱- دانش آموخته دانشگاه تربیت مدرس، دانشکده کشاورزی، گروه علوم و صنایع غذایی

۲- دانشیار دانشگاه تربیت مدرس، دانشکده کشاورزی، گروه علوم و صنایع غذایی

۳- استادیار دانشگاه تربیت مدرس، دانشکده کشاورزی، گروه علوم و صنایع غذایی

چکیده

تأثیر دی اکسید کربن (۰/۵ لیتر بر دقیقه به مدت ۵ دقیقه) و آب پرتقال (۳۰٪ حجمی آب هویج) بر روی مشخصات فیزیکی، شیمیایی و میکروبی آب هویج در طول نگهداری سه ماهه بررسی شد. CO_2 و آب پرتقال باعث کاهش pH آب هویج شدند، لیکن وقتی آب پرتقال (۳۰٪) اضافه شد کاهش pH بیشتر بود. نمونه های مخلوط آب پرتقال و آب هویج اسیدیته بیشتر و pH کمتری داشتند. اختلاط آب پرتقال با آب هویج موجب افزایش رسوب، کاهش در مقادیر a^* و b^* و افزایش L^* شد. نتایج این تحقیق نشان داد که کاهش شکر، درصد رسوب آب هویج پاستوریزه شده را افزایش می دهد. در هیچکدام از تیمارها رشد میکروبی مشاهده نشد.

کلید واژگان: آب هویج-پرتقال، کربناته کردن، فیزیکو شیمیایی، زمان نگهداری، مقدار شکر

* مسئول مکاتبات: hamidy_z@modares.ac.ir