



Optimization of Foaming Parameters and evaluation Foam-Mat Drying of white button mushroom (*Agaricus bisporus*)

Atena Pasban,^{1,2} . Mohebat Mohebbi,^{1*} . Hashem Pour Azarang,¹ . Mehdi Varidi,¹ . Aref Abbasi,¹

1. Department of Food Science and Technology, Faculty of Agriculture, Ferdowsi University of Mashhad, Mashhad, Iran

2. Department of Nutrition, School of Health, North Khorasan University of Medical Sciences, Bojnurd, Iran

ARTICLE INFO

Article History:

Received 2022/ 04/ 03

Accepted 2022/ 07/ 02

Keywords:

Button mushroom,
Foam mat drying,
Modelling,
Physicochemical properties.

DOI: 10.22034/FSCT.19.129.41

DOR: 20.1001.1.20088787.1401.19.129.3.0

*Corresponding Author E-Mail:
m-mohebbi@um.ac.ir

ABSTRACT

Recently, the quality of powder product was considered in food industry. Foam mat drying is an economical alternative to spray, drum and freeze-drying for the production of food powders. The high porosity and surface-to-volume ratio of foams during foam-mat drying improve final product quality. In this work, foaming properties of white button mushroom (*Agaricus bisporus*) were analyzed by response surface methodology. For optimization, xanthan gum (XG) solution range was considered between 0.05–0.3% w/w, mushroom concentration (water: mushroom puree) range was considered 1:0.5– 1:3.5 w/w and whipping time range was considered between 2–8 min. Based on face centered central composite design (CCF), twenty tests were done. Analysis of variance showed that the quadratic model has considerable effects on both responses. In this research, the optimized point was obtained at xanthan gum solution 0.2% w/w, mushroom concentration (water:mushroom puree) = 1:2.2 and whipping time = 6.19 min and the predicted values for foam density and drainage volume were 0.56 g cm⁻³ and 1.8 mL, respectively. At the second step, optimized foam dried at 50, 65 and 80°C and then drying behavior of optimized foam was investigated by different mathematical models. The results indicated that the Midilli model is a high accurate model for evaluating the drying behavior of mushroom foam. At the third step, the influence of the drying temperatures on some qualitative characteristics of foam-mat dried mushroom powder was investigated. It was found, as the temperature increased, water activity and water binding capacity decreased. Also, Lightness parameter for the temperature of 65°C was higher than other temperature. So, white button mushrooms can be processed into powder and used as a functional or nutritional addition in a variety of food products.

1- Introduction

Button mushroom (*Agaricus bisporus*) is the most important variety of edible mushrooms and has high acceptance due to their nutrition facts and health benefits [1]. This mushroom is the most widely planted and consumed mushroom throughout the world and it contributes to about 40% of the world production of mushroom [2]. The high moisture content of these mushrooms has restricted their use by customers. Fresh mushrooms after picking are more perishable and have a short shelf life, so it should be suitable preservation method are considered [3]. Dehydration is one of the effective preservation methods that has been applied worldwide for the preservation of the fresh mushrooms. It should be noted, mushroom structure is very susceptible to heating, so selection of the efficient dehydration techniques more important on quality of the dried mushroom like color, rehydration ability and other properties. In recent years, foam mat drying (FMD) as an efficient method for drying liquid product had been implemented. In this process, food samples are incorporated with foaming agents and subsequently whipped to make stable foam and then dried by dryer. In the next step, dried powder removed from the tray, milled and packaged and is ready to use. High surface area of foam product improves the mass transfer rates and allows operation done at lower temperatures and shorter drying times. It should be noted, the dried product by foam mat drying has a better-quality characteristics like color. it is porous structure so it can be easily reconstituted [4, 5, 6, 7]. This method has been successfully introduced and applied to many commercially successful powder such as tomato juice [8, 9] and tomato pulp [10], Alphonso mango pulp [11], Mango Pulp [12], beetroot [13], papaya pulp [14, 15].

To produce the best quality of powder in FMD method, the porous structure of the foam should be stable for at least one hour, so addition of the stabilizing and foaming agents is essential for the preparation of stable foam [5, 16]. With regard to the previous studies of Aremu et al. [17] and preliminary experiments, proteins in the structure of the white button mushroom are able to make foam, therefore in this study, researchers

conducted foam formation only with foam stabilizers.

Xanthan gum (XG) is an exo-polysaccharides widely used in food process manufacturing. In this study, XG was employed as a foam stabilizer. Considerable increase in the viscosity of liquid samples and high-water solubility in different temperature ranges are some of the characteristics of the xanthan gum [18]. Moreover, XG could prevent the trapping of air during whipping or mechanical mixing which results in a reduction of the foam expansion [19]. Furthermore, Muthukumaran [20] showed xanthan gum provides excellent stability compared to other investigated stabilizers for producing satisfactory egg white foam.

The aim of the present work were: (1) Investigation the influences of whipping time (WT), mushroom concentration (MC), and xanthan gum solution (XG) on the foam density (FD) and foam stability (DV) of mushroom foam and optimization the foaming parameters with response surface methodology (RSM), (2) Investigation the FMD curves and choosing a high accuracy model for prediction of the drying process, (3) Evaluation the effect of the FMD temperatures on some qualitative characteristics of foam mat dried mushroom powder including (water activity (aw), water binding capacity (WBC), and color).

2- Materials and methods

2-1- Material

In this study, the fresh mushroom was prepared from the local markets of Mashhad, Iran. XG (Ceragem Chemical, USA) and sodium metabisulphite (Sigma Aldrich, USA) were bought.

2-2- Sample preparation

2-2-1- Preparation of button mushroom puree

To prepare mushroom puree, at the first step, sliced mushrooms were immersed into sodium metabisulphite 2% (w/w) solution for 10 min. Sodium metabisulphite prevents browning reactions and prevents undesirable discoloration during crushing. Afterward, for getting a homogeneous puree mushroom, sliced mushrooms were crushed by a kitchen blender (Tefal, 210 w) at 15000 rpm for 1 min.

2-2-2- Preparation of gum's solutions

For preparation of XG's solutions, 1 g XG was mixed with 100 ml distilled water and was

homogenized. Afterward, XG solutions was placed in the refrigerator (Hymallia, Iran) about one day to improved hydration.

2-3- Experimental Design and Statistical Analysis

In this research, CCF design with three input variables including, XG (0.05–0.3 w/w), WT (2–8 min) and MC (1:0.5 – 1:3.5 w/w) at three levels was implemented to optimize the foaming condition (FD and DV) of mushroom foam. 20 tests were done based on the CCF design and the center point in the design was repeated six times to evaluate the repeatability of the method. The input variables range, were selected based on the preliminary experiments. The plan of central composite design in coded levels of the three input variables is presented in Table 1. A second order polynomial

equation (Eq. (1)) was fitted to each of the responses:

$$Y_k = \beta_{k0} + \sum_{i=1}^3 \beta_{ki} x_i + \sum_{i=1}^3 \beta_{kii} x_i^2 + \sum_{i=1}^3 \sum_{j=1}^3 \beta_{kij} x_i x_j + \varepsilon_k \quad (1)$$

Where Y_k = response variable (Y_1 = FD; Y_2 = DV) and x_i , and x_j represent the coded input variables ($i=1, 2, 3=X, M, W$), respectively. Where, X = XG solution, M = MC (water:mushroom puree), W = WT. The coefficients of the polynomial are represented by B_{k0} . B_{k0} is the value of the fitted response at the center point of the design, i.e., point (0, 0, 0) and B_{ki} , B_{kii} and B_{kij} which are the linear, quadratic and interaction regression coefficients, respectively.

Table 1 Responses acquired for different process parameter combinations

Experiment No.	Process Variables(coded value)			Response	
	XG (X)	WT (W)	MC (M)	DV (ml)	FD (gcm ⁻³)
1	0.17(0)	8(+1)	1:2(0)	5	0.58
2	0.17(0)	5(0)	1:2(0)	3	0.58
3	0.17(0)	5(0)	1:2(0)	2	0.57
4	0.17(0)	5(0)	1:0.5(-1)	0	0.65
5	0.3(+1)	2(-1)	1:3.5(+1)	1.80	0.8
6	0.05(-1)	8(+1)	1:0.5(-1)	10.50	0.84
7	0.05(-1)	2(-1)	1:0.5(-1)	12	0.91
8	0.3(+1)	5(0)	1:2(0)	0	0.60
9	0.17(0)	5(0)	1:2(0)	4	0.62
10	0.3(+1)	8(+1)	1:3.5(+1)	2	0.64
11	0.3(+1)	8(+1)	1:0.5(-1)	0	0.58
12	0.17(0)	5(0)	1:2(0)	3	0.59
13	0.17(0)	5(0)	1:2(0)	4	0.62
14	0.17(0)	2(-1)	1:2(0)	4.75	0.66
15	0.05(-1)	8(+1)	1:3.5(+1)	25	0.55
16	0.17(0)	5(0)	1:2(0)	1	0.59
17	0.05(-1)	5(0)	1:2(0)	23	0.55
18	0.05(-1)	2(-1)	1:3.5(+1)	33	0.58
19	0.17(0)	5(0)	1:3.5(+1)	4	0.55
20	0.3(+1)	2(-1)	1:0.5(-1)	0	0.77

Regression analysis and analysis of variance (ANOVA) were employed to fit the models to the experimental value and evaluate the statistical significance of the model terms represented by Eq. (1).

Evaluation of the models was done by model analysis, lack-of-fit test, R^2 , adjusted R^2 and predicted R^2 coefficients [21].

2-4- Foam preparation

In this research, the specific amount of stabilizer, mushroom puree and distilled water recommended by CCF design, were mixed and whipped using stirrer (model No. SM88, Sonny, China) with a speed of 1500 rpm (maximum speed) at defined times. A flow chart of the mushroom foam preparation is presented in Fig. 1

2-5- Evaluating the foam Density

For evaluating the FD parameter, the mass of the constant volume of the foam (50ml) was measured at ambient temperature and then foam density was calculated from Eq. 2[3].

$$\text{Foam density} = \frac{\text{Weight of foam (g)}}{\text{Volume of Foam (cm}^3\text{)}} \quad (2)$$

2-6- Evaluating the drainage volume

Drainage volume was evaluated with the method provided by Bag et al.[22] and Sauter and Montoure, [23] with little changes. Thus, the mushroom foam was poured into the Buchner filter (80 mm diameter) that covered with mesh cloth. After one hour, the amount of liquid drainage from the foam sample was considered as a foam stability parameter.

2-7- Drying procedure

The optimized mushroom foams were put uniformly in aluminum plates (9.5 cm diameter) with a foam thickness about 3.0 mm, and then were dried at 50, 65, and 80 °C. Drying experiment was done in laboratory hot air drier (Soroush Medical Company, Khorasan Razavi province) with centrifugal fan and air velocity of 1.5 m/s. Weight loss recorded by digital balance (model AND.EK-300i, Japan) at regular intervals until the mushroom foam samples to get a stable weight. Two replications were carried out for each temperature.




Fig 1 Flowchart for preparation of mushroom foam.

2-8- Data analysis

The statistical analysis of the data, three dimensional (3D) plotting and optimization of foaming parameters of mushroom foams was done by Design Expert 6.0.2 software (Stat Ease Inc., Minneapolis, MN). Numerical and graphical optimization methods in Design Expert software was applied for optimizing the independent variables including MC, XG and WT. All the process parameter selected in their specific range except xanthan gum solution which maximized level was chosen at 0.2%. The desired goals for both responses were at minimized range.

2-9- Mathematical modeling of drying process

Mathematical modeling of drying process of mushroom foams was done by twelve moisture ratio models (Table 2) and high accuracy model for simulation drying process was selected.

The moisture ratio (MR) was simplified to M/M_o instead of the $(M - M_e)/(M_o - M_e)$ which M , M_o , M_e are the moisture content of the foam at each moment, initial moisture

content and equilibrium moisture contents, respectively[25, 26].

Evaluation the mathematical models for simulation the thin layer drying process was done by R^2 , χ^2 , RMSE and SSE values which defined as follows (Eq. 3-6)[24]:

$$R^2 = \frac{\sum_{j=1}^n (MR_{exp,j} - MR_{pre})^2 - \sum_{j=1}^n (MR_{exp,j} - MR_{pre,i})^2}{\sum_{j=1}^n (MR_{exp,j} - MR_{pre})^2} \quad (3)$$

$$\chi^2 = \frac{\sum_{i=1}^N (MR_{exp,i} - MR_{pre,i})^2}{N - n} \quad (4)$$

$$RMSE = \left[\frac{1}{N} \sum_{i=1}^N (MR_{pre,i} - MR_{exp,i})^2 \right]^{1/2} \quad (5)$$

$$SSE = \frac{\sum_{j=1}^n (MR_{pre,i} - MR_{exp,i})^2}{n} \quad (6)$$

Where $MR_{pre,i}$, $MR_{exp,i}$, N and n are the i th predicted MR, the i th experimental MR, the number of observations and the number of constants in drying model, respectively. MATLAB version 7.0 software was applied to examine the constants and coefficients of the mathematical models.

Table 2 Mathematical models references.

Model name	Model	References
Newton	$MR = \exp(-kt)$	O'Callaghan et al. (1971) . ^[27]
Page	$MR = \exp(-kt^n)$	Page (1949) . ^[28]
Modified page	$MR = \exp[(-kt)^n]$	Kingsly and Singh (2007) . ^[29]
Henderson & Pabis	$MR = a \exp(-kt)$	Kingsly and Singh (2007) . ^[30]
Modified Henderson & Pabis	$MR = a \exp(-kt) + b \exp(-gt) + c \exp(-ht)$	Karathanos (1999) . ^[31]
Logarithmic	$MR = a \exp(-kt) + c$	Togrul and Pehlivan (2002) . ^[32]
Appromiximation of diffusion	$MR = a \exp(-kt) + (1 - a) \exp(-kbt)$	Yaldiz et al. (2001) . ^[33]
Simplified Fick's diffusion equation	$MR = a \exp\left(-c \left(\frac{t}{L^2}\right)\right)$	Diamante and Munro (1991) . ^[34]
Midilli	$MR = a \exp(-kt^n) + bt$	Midilli et al. (2002) . ^[35]
Jena & Das	$MR = a \exp(-kt + b\sqrt{t}) + c$	Jena and Das (2007) . ^[36]
Logestic	$MR = \frac{a}{1 + b \exp(kt)}$	Chandra and Singh (1995) . ^[37]
Weibull distribution	$MR = a - b \exp(-kt^n)$	Machado et al. (1998) . ^[38]

2-10- Analysis of the powder

2-10-1- Water activity (aw)

Determination of aw of mushroom powders was evaluated with water activity meter (HygroLab 2, Rotronic Instrument Corp, USA) at constant temperature 24.5 ± 0.5 °C. In each experiment, 2 g of the mushroom powder was transferred in the instrument cup and aw was determined.

2-10-2- Water binding capacity (WBC)

WBC was determined from the technique implemented by McConnell et al. [39] with a little change. In this method, 1 g of mushroom powder samples was suspended in 30mL of distilled water and kept for 18 hours in ambient temperature. Then mixture was centrifuged (Sigma, 2-16KC, Germany) at 3400 g-force for 40 min. The volume of liquid separated after centrifugation (supernatant) was evaluated and WBC was determined by using the below equation (Eq.7):

$$\text{water binding capacity } \left(\frac{\text{ml}}{\text{gr}} \right) = \frac{\text{volume of water added} - \text{volume of supernatant}}{\text{weight of sample}} \quad (7)$$

2-10-3- Color evaluation

The color of the mushroom foams was evaluated with Chroma-meter (Chroma Meter model CR-410, Minolta Co. Ltd., Osaka, Japan) and L^* (lightness) and a^* (redness), parameters were examined. Calibration was done by using a standard white plate. To evaluate the color parameter, foam samples poured in a plate and then color surface was measured by Chroma-meter.

3- Results and discussion

3-1- Evaluating the Foaming properties

The influences of process parameters on foaming characteristics of mushroom foams are presented in Table 1.

From the ANOVA, it was found the quadratic models were highly significant ($p < 0.001$) for two responses (Table 3). The high value of R^2 , adjusted R^2 and predicted R^2 for response variables and non-significant lack of fit indicated that models have high accuracy for predicting response at any combination of variables in experimental range.

The optimum values with desirability of 0.950 was found at XG solution (0.2% w/w), MC (1:2.2) and WT (6.19 min). The predicted values for FD and DV parameters at optimum point were 0.56 gcm^{-3} and 1.8 mL , respectively.

Table 3 Analysis of variance (ANOVA) for response surface models (Eqs. 8 and 9)

Source	Drainage volume			Foam density		
	Sum of squares	F-Value	Prob > F	Sum of squares	F-Value	Prob > F
(a) Model analysis						
Mean	952.89			8.31		
Linear	1189.69	13.21	0.0001	0.066	2.58	0.0896
2Fi	142.33	1.82	0.1925	0.072	4.84	0.0179
Quadratic	301.56	27.48	<0.0001	0.056	21.22	0.0001
Cubic	27.69	4.67	0.0470	0.00229	0.53	0.7203
Residual	8.89			0.00651		
Total	2623.05			8.51		
(b) Lack of fit						
Linear	473.64	31.51	0.0007	0.14	34.80	0.0005
2fi	331.33	30.30	0.0008	0.063	22.28	0.0017
Quadratic	29.74	4.35	0.0662	7.046×10^{-3}	3.98	0.0780
Cubic	2.06	1.50	0.2746	7.748×10^{-3}	13.40	0.0146
Pure error	6.83			1.772×10^{-3}		
(c) R-square analysis	Adjusted R-Squared	Predicted R-Squared	PRESS	Adjusted R-Squared	Predicted R-Squared	PRESS
Linear	0.6584	0.4675	889.33	0.1998	-0.4350	0.29
2fi	0.7041	-0.0830	1808.87	0.5345	-1.4962	0.51
Quadratic	0.9584	0.8084	319.93	0.9178	0.8027	0.040
Cubic	0.9831	-0.5181	2535.47	0.8987	-27.6221	5.84

3-2- Foam density (FD)

FD determination is one of the most common methods for evaluation the foam expansion. As shown in Table 1, FD was in range from 0.55 to 0.91 gcm^{-3} . Also, R^2 and predicted R^2 values were 0.947 and 0.826 that were in good agree with adjusted R^2 of 0.917 (Table 4). In the current study, there were generally significant linear correlation between MC and WT at 0.1%. Also, the relationship between XG solution – MC and XG solution – WT were significant at 0.1% and 5%, respectively. Quadratic effect of MC and WT were significant at 5% and 0.1% levels, respectively, whereas XG solution was not significant at 5% (Table 4).

The effect of input variables on FD parameter was explained by the below equation (Eq. 8):

$$FD = +0.59 - 0.00502X - 0.062M - 0.053W + 0.047M^2 + 0.07W^2 + 0.089XM - 0.031XW \quad (8)$$

Where, X is XG solution/100 g; M is the MC(w/w) and W is WT (min). The response surface plots for FD scores in relation to MC – XG solution and XG solution – WT were represented in Figs. 2 and 3, respectively. As the MC decreased, the FD increased significantly ($p < 0.001$). Similar results were reported for bael fruit [22] and star fruit [4].

High viscosity from increasing MC factor, prevented the incorporating air into foam during whipping and FD decreased significantly [7, 23]. Also, it is observed, in low level of MC parameter, with increasing the XG solution, FD parameter decreased significantly (Fig. 2). This behavior can be explained based on protein–polysaccharide interaction and improving the foaming power. According to Laneuville et al. [41] polysaccharides like XG, through increasing the viscosity of the suspension,

reducing protein – protein interactions, shielding active charged groups and decreasing the collision rate between molecules, prevent the protein aggregation and improving the foaming properties in foam system.

As shown in Fig. 3, FD decreased steadily with increasing WT. It is from, aerating the foam samples during whipping and decreasing the FD. It should be noted that, after the initial decreasing foam density, a little increase in

foam density was observed about 6 min of WT. This maybe occurred due to extended of the WT which make the thinning of the liquid film between the bubbles that cause the bubbles structure to collapse due to the mechanical force from stirrer [40, 42]. Also, increasing the whipping time or “overbeating” make to accelerate aggregation and precipitation in proteins structure of foam samples and decreasing the foaming capacity of mushroom foam, consequently [43].

Table 4 Regression summary and ANOVA table for Foam density

Source	Sum of squares	DF	Mean squares	F-Value	Prob > F
Model	0.19	7	0.028	31.08	< 0.0001 significant
X	2.520×10^{-4}	1	2.52×10^{-4}	0.28	0.6040
M	0.038	1	0.038	43	< 0.0001
W	0.028	1	0.028	31.58	0.0001
M ²	6.968×10^{-3}	1	6.968×10^{-3}	7.84	0.0160
W ²	0.016	1	0.016	17.79	0.0012
XM	0.064	1	0.064	72.11	< 0.0001
XW	7.454×10^{-3}		7.454×10^{-3}	8.39	0.0134
Residual	0.011	12	8.882×10^{-4}	-----	-----
Lack-of-fit	8.887×10^{-3}	7	1.270×10^{-3}	3.58	0.0895 not significant
Pure error	1.772×10^{-3}	5	3.543×10^{-4}	-----	
Total	0.2	19			
R-Squared	0.9477				
Adj R-Squared	0.9172				
Pred R-Squared	0.8265				




Fig 2 Effect of input variables (MC and XG solutions) on FD parameters of mushroom foams.




Fig 3 Effect of input variables (XG solution and WT) on FD parameters of mushroom foams.

3-3- Drainage volume

Foam structure naturally are unstable and delicate. High surface energy in the air – water interface and the different density between two phases makes foams thermodynamically unstable in nature[20].

Foam stability commonly evaluated by DV parameter, which is related the flow of the liquid phase through foam capillary or external forces[44].

In the current study, there were significant linear correlation between XG solution and MC at the 0.1 % level of significance. Also, the quadratic effect of XG solution was found to be significant at the same level. There was a significant interaction between XG solution and MC at 0.1% level (Table 5).

The effect of input variables on DV parameter was explained by the below equation (Eq. 9):

$$DV = +3.08 - 9.97X + 4.33M + 7.66X^2 - 3.96XM \quad (9)$$

Table 5 Regression summary and ANOVA table for Drainage volume

Source	Sum of squares	DF	Mean squares	F-Value	Prob > F
Model	1600.1	4	400.03	85.65	< 0.0001 significant
X	994.01	1	994.01	212.83	< 0.0001
M	187.49	1	187.49	40.14	< 0.0001
X ²	293	1	293	62.73	< 0.0001
XM	125.61	1	125.61	26.89	0.0001
Residual	70.06	15	4.67	-----	-----
Lack-of-fit	63.22	10	6.32	4.63	0.0524 not significant
Pure error	6.83	5	1.37	-----	-----
Total	1670.16	9			
R-Squared	0.9581				
Adj R-Squared	0.9469				
Pred R-Squared	0.9045				

Where, X is the XG solution/g/100 g of the mixture; M is the MC(w/w) and W is the WT (min). The variations in DV parameter of the mushroom foams with XG solution and MC in 3 D surface plot are presented in Fig. 4. It is clear that XG solution had more effect on DV. As the XG solution increased, continuous decrease in DV parameter was occurred. XG by increasing the viscosity of liquid foam, improved the interfacial viscoelasticity of foam lamellae and helps to stabilize the foam system [45, 46]. Our results are in agreement with those of Azizpour et al. [6], Abbasi et al. [7] and Salahi et al. [19].

As the plot shows, with decreasing the MC factor (water: mushroom puree) or in other word with decreasing the dilution rate, DV parameter decreased significantly. Enhancement of foam stability could be from increasing the total solid content in mushroom suspensions specially protein compounds which improve the water holding in foam structure and improve foam stability. According to German & Phillips [47] and Wilde & Clark [48], proteins structures have a

major role in increasing the foam stability by reducing the interfacial tension to low level and making more mechanical strength in the bubble walls by interacting with lamella and form the electrostatic, hydrophobic, and covalent and hydrogen bonding.




Fig 4 Effect of input variables (MC and XG solution) on DV parameters of mushroom foams.

3-4- Evaluation of drying process

Evaluation of FMD curves of optimized foams at FMD temperatures 50, 65 and 80 °C was shown in Fig. 5. It is found that by increasing

the FMD temperature, the FMD time decreased consequently, and FMD temperature has a significant influence on the FMD time. For example, when the FMD temperature was increased from 50 to 80 °C, the FMD time decreased approximately 50% which related to the increasing of the heat and mass transfer rate during the foam samples. FMD time required to reach the equilibrium moisture content were about 87, 57 and 39 min at 50, 65 and 80 °C, respectively. Similar results for FMD of banana and shrimp foam samples have been reported [49, 50].




Fig 5 drying curves for drying temperatures.

3-5- Fitting of the FMD curves

The MR of optimized foam at three FMD temperatures (50, 65 and 80 °C) was fitted in

twelve drying models and the statistical analysis results are presented in Table 6. In all drying models, R^2 values were bigger than 0.93, which indicates that all mathematical models have high correlation with experimental values. As shown in Table 6, Midilli model has the maximum value of R^2 and minimum values of χ^2 , SSE and RMSE. In Fig 6, the MR values from Midilli model were compared by experimental values, which confirm Midilli model is a high accurate model for evaluating FMD behavior of mushroom foams. Constants and coefficients of the Midilli model at FMD temperature are presented in Table 7.




Fig 6 Experimental and predicted MR values for different drying conditions.

Table 6 Statistical results of 12 models at three drying temperatures

Drying air temperature												NO
80				65				50				
SSE	RMSE	X ²	R ²	SSE	RMSE	X ²	R ²	SSE	RMSE	X ²	R ²	
0.1173	0.0884	0.00733	0.936	0.1918	0.08759	0.00737	0.935	0.0222	0.0868	0.00742	0.935	1
0.0154	0.0332	0.00096	0.991	0.0279	0.03414	0.00107	0.990	0.0194	0.0258	0.00062	0.994	2
0.0154	0.0345	0.00012	0.991	0.0417	0.04261	0.00189	0.989	0.0201	0.0268	0.00074	0.994	3
0.0876	0.0791	0.00626	0.952	0.148	0.07854	0.00643	0.950	0.1572	0.0736	0.00561	0.954	4
0.0788	0.0887	0.00754	0.957	0.1825	0.09554	0.00668	0.983	0.0283	0.0336	0.00118	0.991	5
0.0183	0.0375	0.00141	0.990	0.0580	0.05025	0.00264	0.980	0.0354	0.0351	0.00127	0.990	6
0.0239	0.0428	0.00184	0.987	0.0491	0.04621	0.00225	0.983	0.0391	0.0373	0.00145	0.988	7
0.0876	0.0821	0.00674	0.952	0.148	0.08023	0.00672	0.950	0.1572	0.0749	0.00582	0.954	8
0.0074	0.0249	0.00062	0.995	0.0098	0.02118	0.00047	0.996	0.0038	0.0119	0.00019	0.998	9
0.0110	0.0303	0.00091	0.994	0.0391	0.0422	0.00187	0.986	0.0191	0.0266	0.00073	0.994	10
0.0876	0.0821	0.00675	0.952	0.148	0.08023	0.00673	0.950	0.1572	0.0749	0.00582	0.954	11
0.0068	0.0268	0.00071	0.995	0.0156	0.02665	0.00074	0.994	0.0069	0.0160	0.00028	0.998	12

Table 7 Statistical results of Midilli model and its constants and regression coefficients at three drying temperatures

Drying air temperature (°C)	a	B	k	n	SSE	RSME	X ²	R ²
50	0.9869	-0.04655	-0.05565	0.7237	0.0038	0.0119	0.00019	0.998
65	0.9862	-0.06049	-0.06322	0.7382	0.0098	0.02118	0.00047	0.996
80	0.9984	-0.08422	-0.07814	0.7448	0.0074	0.0249	0.00062	0.995

3-6- Evaluating the mushroom powder

3-6-1- Water activity

The aw values of mushroom powder are presented in Table 8. It was found as the FMD temperature increased, the aw of mushroom powders decreased significantly ($P < 0.05$). It should be noted, in all mushroom powder samples, aw was below the 0.6, that indicate the available water for chemical reactions and microbiological growth is insufficient and mushroom powder from FMD will have a good shelf life [52]. A considerable reduction in aw parameter could be from the porous structure of mushroom foams which accelerating the moisture loss during drying process. Also unfolding and denaturation of proteins structure make the holding water in protein structure reduced and consequently aw reduced [42]. Some research in their studies on FMD of foods reported the same conclusion [52, 53].

3-6-2- Color

The most important factor that is expected to affect marketability of the food dried product is color. Millard reaction and browning enzymatic are the most important reasons for color changing in mushroom during drying process. According to the previous studies, the color parameters L^* and a^* , are well correlated to color changes and darkening due to browning reactions in food product [54]. By increasing the browning reaction L^* parameter will decrease and a^* parameter will increase. Therefore, in present study L^* and a^* parameters were considered as color evaluation index (Table 8).

As shown in Table 8, L^* for mushroom powders dried the 50 and 80°C were higher than the mushroom powder which dried in

65°C. Also, a^* in mushroom powder dried at 65°C was more than the mushroom powder dried in 50 and 80°C. In other word, mushroom powder from FMD at 65°C has less lightness (more browning) compared to the dried powder produced in 50 and 80°C. Whereas, there was not significant difference between L^* parameters for mushroom powders dried at 50 and 80°C. It should be noted that, temperature and time of drying process are the important parameter on color change and browning reaction during drying process [55]. This result can be from the less browning reaction (especially enzymatic browning) at 50°C compared with the drying temperature of 65°C. Also, longer drying time at 65°C caused the final powders has lower lightness than the mushroom powder dried at 80°C. In other word low drying time in drying process at 80 makes the final powder has a better color.

3-6-3- Water binding capacity

Another important parameter for evaluation the quality of the food dried products is water binding capacity (WBC). WBC represents the amount of water that dried product can be absorb, which depends on the structure of proteins and the capability to hydration [56].

As shown in Table 8 by increasing the drying temperature, WBC decreased, which is related to the denaturation of proteins that causes the aggregation and coagulation of the protein molecules [52].

It should be noted, reduction in WBC with increasing drying temperature was insignificant, that it could be from the presence of xanthan gum that has heat resistance property and improve WBC parameter.

Table 8 Effect of drying temperature on some properties of mushroom powder.

Water binding capacity	Color		Water activity	Drying temperature (°C)
	a^*	L^*		
4/23 ^a	1/20 ± 0 ^a 728	68/44 ± 0 ^a 077	0/142 ± 0 ^a 008	50
4/085 ^a	4/17 ± 0 ^b 721	59/89 ± 0 ^b 756	0/079 ± 0 ^b 026	65
3/39 ^a	2/18 ± 0 ^{ab} 474	68/19 ± 0 ^a 848	0/050 ± 0/013 ^b	80

Different letters (a–c) within a same column differ significantly ($P < 0.05$)

4- Conclusion

In this paper, RSM was implemented for determination of the optimized point of foaming condition of mushroom foam with XG solution in the range of 0.05–0.3% (w/w), MC of 1/0.5–1/3.5 (w/w) and WT of 2–8 min as input variables. The results showed, the developed model were more efficient tools and techniques for predicting foaming properties of mushroom foam to the experimental data. The optimized foaming conditions point was obtained at XG = 0.2% w/w, MC = 1:2.2 (w/w) and WT 6.19 min. Drying curve studies showed that FMD temperature has a considerable influence on the diffusion of water through the foam, and by increasing the FMD temperatures, moisture lost was increased. Evaluating the mathematical modeling the FMD curves was shown that Midilli model is appropriate and high accurate model for investigating the FMD behavior of mushroom foams. Evaluation the physicochemical properties of mushroom powder show, with increasing the FMD temperature, water activity and water binding capacity decreased consequently. also, Lightness parameter for the temperature of 50, 65 and 80°C was bigger than the 65°C.

5- References

- [1] E. V. Crisan and A. Sands, Nutritional Value of Edible Mushroom, In: S. T. Chang and W. A. Hayer, Eds., *Biology and Cultivation of Edible Mushrooms*, Academic Press, New York, 137-168 (1978)
- [2] I. Doymaz, Drying kinetics and rehydration characteristics of convective hot-air dried white button mushroom slices. *J. Chem.* (2014)
- [3] M. Brenan, G. Le port, A. Pulvirenti, and R. Gormley, (2000) <https://doi.org/10.1006/fstl.2000.0657>
- [4] A.A. Karim, and C.C. Wai, Characteristics of foam prepared from starfruit (*Averrhoa carambola* L.) puree by using methyl cellulose. *J. Food. Hydrocoll.* 13, 203–210 (1999)
- [5] D.M. Kadam, R.T. Patil, and P. Kaushik, Foam Mat Drying of Fruit and Vegetable Products. In *Drying of Foods, Vegetables and Fruits*, (Jangam, S.V., Law, C.L. & Mujumdar, A.S. eds), 113–124, Central Institute of Post Harvest Eng. Technol. Ludhiana, India. (2010)
- [6] M. Azizpour, M. Mohebbi, M. H. Haddad Khodaparast, & M. Varidi, Optimization of foaming parameters and investigating the effects of drying temperature on the foam-mat drying of shrimp (*Penaeus indicus*). *J. Dry. Technol.* 32, 374–384. (2014).
- [7] E. Abbasi, M. Azizpour, Evaluation of physicochemical properties of foam mat dried sour cherry powder. *LWT Food Sci. Technol.* 68, 105–10 (2016)
- [8] D. M. Kadam, S. Balasubramanian, S. (2011). Foam mat drying of tomato juice. *J. food processing and preservation*, 35, 488–495.
- [9] M. A. Hossain, S. Mitra, M. Belal, W. Zaman, (2021). Effect of foaming agent concentration and drying temperature on biochemical properties of foam mat dried tomato powder. *Food Research*, 5, 291–297.
- [10] O. S. Qadri, A. K. Srivastava, (2014). Effect of microwave power on foam-mat drying of tomato pulp. *Agricultural Engineering International: CIGR Journal*, 16, 238–244.
- [11] P. Rajkumar, R. Kailappan, R. Viswanathan, G. S. V. Raghavan, C. Ratti, C. Foam mat drying of Alphonso mango pulp. *J. Dry. Technol.* 25, 357–365. (2007).
- [12] A. M. Chaux-Gutiérrez, A. B. Santos, D. M. Granda-Restrepo, M. A. Mauro, Foam mat drying of mango: Effect of processing parameters on the drying kinetic and product quality. *Drying Technology*, 35, 631–641. (2017).
- [13] M. L. Ng, R. Sulaiman, R. Development of beetroot (*Beta vulgaris*) powder using foam mat drying. *Lwt*, 88, 80–86. (2018).
- [14] P. Kandasamy, N. Varadharaju, S. Kalemullah, D. Maladhi, Optimization of process parameters for foam-mat drying of papaya pulp. *J. food Sci. technol*, 51, 2526–2534 (2014).
- [15] A. El-Salam, E. Abd El-Salam, A. M. Ali, K. S. Hammad, Foaming process optimization, drying kinetics and quality of foam mat dried papaya pulp. *J. Food Sci. Tech*, 58, 1449–1461 (2021).
- [16] C. Ratti, and T. Kurda, Drying of foamed biological materials: opportunities and challenges. *J. Dry. Technol.* 24, 1101–1108 (2006)
- [17] M.O. Aremu, S.K. Basu, S.D. Gyar, A. Goyal, P.K. Bhowmik, S. Datta Banik,

- Proximate composition and functional properties of mushroom flours from *Ganoderma* spp., *Omphalotusolearius* (DC.) Sing. & *Hebelomamesophaeum* (Pers.) Quél, Used in Nasarawa State, Nigeria. *Malaysian J. Nutr.* 15, 233–241 (2009)
- [18] J. N. Bemiller, R.L. Whistler, Carbohydrates, in *Food Chemistry*, 3rd ed. O.R. Fennema, Marcel Dekker. New York. 157-233 (1996)
- [19] M. R. Salahi, M. Mohebbi, M. Taghizadeh, Foam - Mat Drying of Cantaloupe (*Cucumis melo*): Optimization of Foaming Parameters and Investigating Drying Characteristics. *J. food processing and preservation*, 39, 1798-1808 (2015)
- [20] A. Muthukumaran, Foam-mat freeze drying of egg white and mathematical modeling. MSc Thesis, In Department of Bio resource Engineering, McGill University (2007).
- [21] J. Lee, L. Ye, W. Landen, R.R. Eitenmiller, Optimization of extraction procedure for the quantification of vitamin E in tomato and broccoli using response surface methodology. *J. Compo. Analysis*, 13, 45–57 (2000)
- [22] S. K. Bag, P.P. Srivastav, H.N. Mishra, Optimization of Process Parameters for Foaming of Bael (*Aegle marmelos* L.) Fruit Pulp. *Food. Bioproc. Technol.* 4, 1450-1458(2011)
- [23] E. A. Sauter, and J.E. Montoure, The relationship of lysosyme content of egg white to volume and stability of foams. *J. Food Sci.* 37, 918–920 (1972)
- [24] N.J. Thakor, S. Sokhansanj, F.W. Sosulski, S. Yannacopoulos, Mass and dimensional changes of single canola kernels during drying. *J. Food. Eng.* 40, 153–160 (1999)
- [25] Z. Wang, J. Sun, X. Liao, F. Chen, G. Zhao, J. Wu, X. Hu, Mathematical modeling on hot air drying of thin layer apple pomace. *Food Res. Int.* 40, pp 39–46 (2007)
- [26] S. C. Chapra, R.P. Canale, Numerical methods for engineers. New York: McGraw-Hill Book Company, 337(1989)
- [27] J. R. O'Callaghan, D.J. Menzies, P.H. Bailey, Digital simulation of agricultural dryer performance. *J. Agric. Eng. Res.* 16(3), 223–244 (1971)
- [28] G. E. Page, Factors influencing the maximum rates of air drying shelled corn in thin layers. MSc thesis. Department of Mechanical Engineering, Purdue University, Purdue, USA (1949)
- [29] G. M. White, I.J. Ross, R. Ponekert, Fully exposed drying of popcorn. *Transactions of the ASAE.* 24, 466–468 (1981)
- [30] S. M. Henderson, and S. Pabis, Grain drying theory. II. Temperature effects on drying coefficients. *J. Agric. Eng. Research.* 6, 169–174 (1961)
- [31] V. T. Karathanos, Determination of water content of dried fruits by drying kinetics. *J. Food. Eng.* 39(4), 337–344 (1999)
- [32] I. T. Togrul, and D. Pehlivan, Mathematical modeling of solar drying of apricots in thin layers. *J. Food. Eng.* 55, 209–216 (2002)
- [33] O. Yaldiz, C. Ertekin, and H.I. Uzun, Mathematical modeling of thin layer solar drying of sultana grapes. *Energy J.* 26(5), 457–465(2001)
- [34] L. M. Diamante, and P.A. Munro, Mathematical modeling of hot air drying of sweet potato slices. *International J. Food.Sci.Technol.* 26(1), 99–109 (1991)
- [35] A. Midilli, H. Kucuk, Z. Yapar, A new model for single layer drying. *J. Drying Technol.* 20(7), 1503–1513 (2002)
- [36] S. Jena, and H. Das, 2007. Modeling for Vacuum Drying Characteristics of Coconut Presscake. *J. Food. Eng.* 79(1), 92–99.
- [37] P. K. Chandra, and R.P. Singh, Applied numerical methods for food and agricultural engineers. Boca Raton, FL: CRC Press. 163–167 (1995)
- [38] M. F. Machado, F.A.R. Oliveira, V. Gekas, and P. Singh, Kinetics of moisture uptake and soluble solids loss by puffed breakfast cereals immersed in water. *International J. Food Sci. Technol.* 33(3), 225–237 (1998)
- [39] A. A. McConnell, M.A. Eastwood, and, W.D. Mitchell, Physical characteristics of vegetable food stuffs", *J. Sci. Food. Agric.* 25, 1457-64 (1974)
- [40] J. J. Bikerman, *Foams*. Springer Verlag, New York (1973)
- [41] S. I. Laneville, P. Paquin, & S.L. Turgeon, Effect of preparation conditions on the characteristics of whey protein-xanthan gum complexes. *J. Food. Hydr.* 14, 305–314 (2000)
- [42] K. O. Falade, K.I. Adeyanju, P.I. Uzo-Peters, Foam-mat drying of cowpea (*Vigna unguiculata*) using

- glycerylmonostearate and egg albumin as foaming agents. *Euro. Food Res. Technol.* 217, 486–491(2003)
- [43] S. Damodaran, Amino Acids, Peptides, and Proteins. In *Food Chemistry*, (Fennema, O.R. ed). 3rd ed. New York, 322–416 (1996)
- [44] W. Drenckhan, S. Langevin, Monodisperse foams in one to three dimensions. *Current Opinion in Colloid and Interface Science*, 15, 341–358 (2010)
- [45] J. P. Heller, M.S. Kuntanukula, Critical review of the foam rheology literature. *Indust. Eng. Chem. Res.* 26, 318–325(1987)
- [46] E. M. Papalamprou, E.A. Makri, V.D. Kiosseoglou, G.I. Doxastakis, Effect of medium molecular weight xanthan gum in rheology and stability of oil in water emulsion stabilized with legume proteins. *J. Sci. Food. Agric.* 85. 1967–1973 (2005)
- [47] J. B. German, L. Phillips, Protein Interactions in Foams: Protein-Gas Phase Interactions in Protein Functionality in Food Systems, N. S. H. G. R. Ziegler, Marcel Dekker Inc, 181–208 (1994).
- [48] P. J. Wilde, D.C. Clark, Foam formation and stability. *Methods of testing protein functionality*. GM Hall, Blackie Academic and Professional, 111–152 (1996)
- [49] R. Thuwapanichayanan, S. Prachayawarakorn, S. Soponronnarit, Drying characteristics and quality of banana foam mat. *J. Food. Eng.* 86, 573–583 (2008)
- [50] S. Thankitsunthorn, C. Thawornphiphatdit, N. Laohaprasit, G. Srzednicki, Effects of drying temperature on quality of dried Indian Gooseberry powder. *Int. Food. Res. J.* 16, 355–361(2009)
- [51] O. R. Fennema, *Food chemistry*, 3rd ed. Marcel Dekker Inc, New York. Goodwin, T.W. (1984). *The biochemistry of the carotenoids* (2nd ed.). London: Chapman and Hall (1996)
- [52] T.S. Franco, C.A. Perussello, L.N. Ellendersen, M.L. Masson, Effects of foam mat drying on physicochemical and microstructural properties of yacon juice powder. *LWT Food Sci. Technol.* 66, 503–513 (2016)
- [53] I. Mandala, E. Anagnostaras, C. Oikonomou, Influence of osmotic dehydration conditions on apple air-drying kinetics and their quality characteristics. *J. Food. Eng.* 69(3), 307–316 (2005)
- [54] H. Qing-guo, Z. Min, A.S. Mujumdar, D. Wei-hua, S. Jin-cai, Effects of different drying methods on the quality changes of granular edamame. *J. Dry. Technol.* 24(8), 1025–1032(2006)
- [55] G. V. Barbosa-Canovas, E. Ortega-Rivas, P. Juliano, H. Yan, *Food powders: Physical properties, processing and functionality* (1st ed.). New York: Kluwer Academic/ Plenum Publishers (2005)
- [56] Zhang, Y., Sharan, S., Rinnan, Å., & Orlén, V. (2021). Survey on Methods for Investigating Protein Functionality and Related Molecular Characteristics. *Foods*, 10(11), 2848.



بهینه سازی شرایط تولید کف و ارزیابی فرآیند خشک کردن کف پوشی قارچ دکمه ای

سفید (*Agaricusbisporus*)آتنا پاسبان^{۱*}، محبت محبی^{۱*}، هاشم پوراآذرننگ^۱، مهدی وریدی^۱، آرزو عباسی^۱

۱- گروه علوم و صنایع غذایی، دانشکده کشاورزی، دانشگاه فردوسی مشهد، مشهد، ایران.

۲- گروه علوم تغذیه، دانشکده بهداشت، دانشگاه علوم پزشکی خراسان شمالی، بجنورد، ایران.

چکیده

اطلاعات مقاله

تاریخ های مقاله :

تاریخ دریافت: ۱۴۰۱/۰۱/۱۴

تاریخ پذیرش: ۱۴۰۱/۰۴/۱۱

کلمات کلیدی:

قارچ دکمه ای،

خشک کردن کف پوشی،

مدلسازی،

خواص فیزیکوشیمیایی.

اخیرا کیفیت فرآورده های پودری در صنایع غذایی مورد توجه قرار گرفته است. خشک کردن به روش کف پوشی یک روش اقتصادی جایگزین برای خشک کردن به روش پاششی، غلطکی و انجمادی برای تولید پودرهای غذایی است. تخلخل و نسبت سطح به حجم بالا در نمونه های کف در حین خشک شدن کف پوشی، کیفیت محصول نهایی را بهبود می بخشد. در این پژوهش، خصوصیات کف زایی قارچ دکمه ای سفید (*Agaricusbisporus*) با روش سطح پاسخ مورد تجزیه و تحلیل قرار گرفت. جهت بهینه سازی، محلول صمغ زانتان (XG) بین ۰/۰۵-۰/۳ درصد وزنی، غلظت سوسپانسیون (آب: پوره قارچ) بین ۰/۵-۱:۳/۵ وزنی/وزنی و محدوده زمان هم زدن بین ۲ تا ۸ دقیقه در نظر گرفته شده است. بر اساس طرح مرکب مرکزی متمرکز شده (CCD)، بیست آزمایش انجام شد. تجزیه و تحلیل واریانس نشان داد که مدل های درجه دوم اثرات قابل توجهی بر هر دو پاسخ دارند. در این پژوهش، نقطه بهینه در محلول صمغ زانتان ۰/۲٪ وزنی/وزنی، غلظت سوسپانسیون (آب: پوره قارچ) ۲/۲: ۱ و زمان هم زدن ۶/۱۹ دقیقه و مقادیر پیش بینی شده برای دانسیته کف و میزان مایع جدا شده از کف به ترتیب ۰/۵۶ گرم در سانتی متر و ۱/۸ میلیلیتر به دست آمد. در مرحله دوم، کف بهینه شده در دماهای ۵۰، ۶۵ و ۸۰ درجه سانتیگراد خشک شد و سپس رفتار خشک شدن کف بهینه شده توسط مدل های مختلف ریاضی بررسی شد. نتایج نشان داد که مدل میدیلی مدلی با دقت بالا برای ارزیابی رفتار خشک کردن کف قارچ است. در مرحله سوم، تأثیر دمای خشک کردن بر برخی ویژگی های کیفی پودر قارچ خشک شده مورد بررسی قرار گرفت. نتایج نشان داد که با افزایش دما، فعالیت آبی و قابلیت جذب آب کاهش یافت. همچنین پارامتر روشنایی در دمای ۶۵ درجه سانتیگراد بالاتر از سایر دماهای خشک کردن بود. بنابراین قارچ دکمه ای سفید را می توان به پودر تبدیل کرد و به عنوان یک افزودنی کاربردی یا تغذیه ای در انواع محصولات غذایی استفاده کرد.

DOI: 10.22034/FSCT.19.129.41
DOR: 20.1001.1.20088787.1401.19.129.3.0

* مسئول مکاتبات:

m-mohebbi@um.ac.ir