

Journal of Food Science and Technology (Iran)

Homepage:www.fsct.modares.ir

Scientific Research

Effect of traditional sourdough on quality characteristics of loaf bread containing sprouted wheat and garlic extract

Sara Safari, Alireza Sadeghi*, Ali Moayedi, Hossein Purabdolah, Alireza Sadeghi Mahoonak

Department of Food Science and Technology, Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran.

ARTICLE INFO	ABSTRACT		
	Application of plant extracts in the formulation of sourdough		
Article History:	can improve sensory characteristics and shelf-life of the		
Received:2024/7/17	produced bread. In the present study, effects of combined		
A 10004/11/10	application of fermented sprouted wheat and garlic extract in the		
Accepted:2024/11/10	formulation of traditional sourdough containing yoghurt and		
Kouwonda	vinegar were investigated on the quality characteristics of loaf		
Keywords:	wheat bread. Based on the results of texture analysis, among the		
garlic extract,	produced breads the lowest crumb hardness was observed in the		
fermented sprouted wheat,	sample containing fermented sprouted wheat along with garlic		
raditional sourdough, extract (2.94 N), which was significantly (p<0.05) lower the			
moldiness.	those of the control sample. In addition, combined application		
	of fermentation and sprouting increased the specific volume and		
	reduced the gumminess of the product. Moreover, although the		
	lowest surface growth of Aspergillus flavus was observed on		
DOL 10 22024/ECOT 22 150 122	breads containing yoghurt-vinegar and also garlic extract alone,		
DOI: 10.22034/FSC1.22.159.133.	there was no significant difference between the fungal growth		
*Corresponding Author E-	rate on the sample containing fermented sprouted wheat along		
sadeghi.gau@gmail.com	with garlic extract and bread containing calcium propionate.		
	Bread containing termented sprouted wheat and control sample		
	had no significant difference in terms of overall acceptability.		
	Accordingly, sprouted wheat sourdough containing garlic		
	extract can be used as a biological preservative to improve the		
	quality characteristics and shelf-life of wheat bread.		

1-Introduction

Bread the production is one of oldest technologies known to humankind. Depending on cultural habits. bread is consumed in various types and forms around the world as a staple food. Wheat bread spoilage can occur due to various factors including bacterial and fungal contamination, with fungal spoilage being the primary [1,2]. concern One approach to controlling bread spoilage is the use of chemical preservatives such as propionic, sorbic, acetic acids. and their salts. which can slow down fungal spoilage. Although these compounds are classified as safe, their use has led to consumer dissatisfaction due to of development some resistant fungi [3,4]. An alternative to chemical preservatives is the use natural of compounds that can reduce eliminate microbial or populations while improving the quality of the produced product. These compounds can also serve natural antifungal additives as to fungal growth and prevent extend shelf-life, thus the bread's reducing public health risks. Among the most important biological preservatives bread in processing fermented grains are plant Several studies and extracts. have shown that plant extracts bioactive contain various components that control can fungal growth due to their phenolic compounds [5,6].

Studies have been conducted on using controlled fermentations as biological preservatives in bread. For example, the effect of controlled fermentation of oat containing Pediococcus pentosaceus starter culture on fungal growth inhibition, texture characteristics, and overall acceptability of wheat bread was evaluated, showing that controlled fermentation can improve these attributes [7]. Another study investigated the impact of controlled fermentation of acorns using Pediococcus acidilactici on the texture and antifungal activity in wheat bread. The results showed that this bacterium positively affected texture and reduced Aspergillus flavus growth on the surface of the produced bread [8]. Additionally, the effect of controlled fermentation of sprouted barley sourdough [9], amaranth [10], sprouted mung beans with ginger extract [11], sprouted clover [12], mung beans [13], quinoa [14], wheat bran [15], wheat germ [16], rice bran [17], and whole wheat flour [18] on technological-functionality, nutritional, and shelf-life characteristics of the produced bread has been confirmed.

There are also reports on using essential oils and plant extracts as biological preservatives in bread production. For instance, the antifungal effect of thyme essential oil on Penicillium paneum in partially baked bread was investigated. The results showed that this essential oil was able to inhibit fungal growth in the bread for up to 45 days [1]. Other researchers combined lactic acid bacteria fermentation with hop extract to improve antifungal properties and create a natural preservative in bread. These studies showed significant antifungal activity of hop extract, delaying fungal growth for up to 14 days [19]. The antifungal activity of essential oils from cinnamon, clove, pepper, camphor, star anise, lemon, and peppermint on three indicator fungi was also investigated, with cinnamon oil showing the greatest effect on all studied fungi [20].

Based on the literature reviewed, there has been no report on the combined use of garlic extract and sprouted wheat in sourdough wheat bread. Thus, the aim of this study was to evaluate the impact of traditional sourdough containing sprouted wheat and garlic extract on the textural, shelf-life, and overall acceptability characteristics of wheat bread.

2-Materials and Methods

2.1. Preparation of raw materials and their evaluation

The characteristics of wheat flour. including moisture, ash, protein, and fat contents, were determined using standard from AACC [21]. methods The carbohydrate percentage was calculated using the formula: 100 - (fat + protein + ash+ moisture). Accordingly, the wheat flour used in this study had 12.4% protein, 0.4% ash, 14.3% moisture, 1.1% fat, and 71.8% carbohydrates. Microbiological culture media and chemical reagents were obtained from reputable commercial brands for laboratory use.

2.2. Preparation of sprouted wheat flour

For sprouting wheat grains, the grains were initially disinfected using 1.25% sodium hypochlorite and then rinsed after 30 minutes [22]. Subsequently, the disinfected grains were hydrated in the dark at 25 °C with 12-hour intervals. The sprouted grains were then dried at 50 °C in an oven (Binder, Germany) for 25 h, and after grinding, sprouted wheat flour was prepared.

2.3. Fermentation using traditional sourdough

The sourdough used in this study was prepared from a mixture of sprouted wheat flour (68%), yogurt (3.5%), vinegar (1.5%), and water (27%) according to the Iranian

standard [23] with minor national modifications. After 24 h of fermentation at 37 °C, the mixture was added to a mix of water and wheat flour (60% water absorption) and baker's yeast (2% w/w). The dough was then fermented for 2 h at 37 before baking. The °C sourdough containing garlic aqueous extract was prepared using the method described by Irakli et al. [24], with the optimal garlic amount determined extract for fermentation. The optimal amount of garlic extract in the bread formulation was determined in the range of 5 to 20% based on overall acceptability scores.

2.4. Production of leavened wheat bread The control sample (containing wheat flour, water, and Saccharomyces cerevisiae) was fermented for 2 h at 37 °C and then baked in an electric oven (Feller, Germany) at 180 °C for 20 minutes. Bread samples containing calcium propionate (0.3% w/w), sprouted wheat, garlic extract, yogurt and vinegar mixture, sourdough (sprouted wheat, yogurt, and vinegar), and sourdough with garlic extract were processed similarly to the control sample. To determine the optimal sourdough amount in the wheat bread formulation, sourdough amounts ranging from 5 to 20% were used, and the optimal amount was determined based on overall acceptability and texture analysis results [25].

2.5. Evaluation of quality characteristics of produced bread

2.5.1. Crumb hardness and specific volume measurement

Hardness of the bread crumb was measured using a texture analyzer (Stable Micro System, UK) with a cylindrical probe at a speed of 50 mm/min and a force of 500 N to achieve 50% compression from the initial height of the sample. The crumb hardness was determined one hour after baking using texture profile analysis [25]. The specific volume of the bread was measured using the rapeseed displacement method [17].

2.5.2. Surface mold growth assessment

To assess the expansion of *A. flavus* on the surface of the produced bread samples, 3 μ L of fungal spore suspension (10⁶ spores/mL) were inoculated onto a sterile paper disk placed at the center of the bread. The samples were incubated for one week at 25 °C. Mold surface growth was compared to the control sample over time by daily photography and analysis using Image J software [26].

2.5.3. Overall acceptability of the produced bread

Sensory evaluation of the produced bread samples was conducted by trained evaluators using a 5-point hedonic scale. This evaluation included assessing the crust color, taste, chewiness, bread aroma, and mouth-feel. Scores were reported from very undesirable (1) to very desirable (5). Overall acceptability of the bread samples was calculated as the average score of these attributes [27].

2.6. Statistical analysis of results

The results were analyzed using a completely randomized design with oneway ANOVA. Mean comparisons were performed using the Least Significant Difference (LSD) test at p<0.05. All tests were conducted in triplicate, and data analysis was performed using SPSS software (version 16) and Microsoft Office Excel 2013 for plotting graphs.

3-Results and Discussion

3.1. Selection of optimal sourdough formulation

Initially, adding more than 20% sourdough led to excessive stickiness in the bread's texture. Therefore, lower percentages of sourdough were used to achieve desirable textural and sensory characteristics in the produced bread. As shown in Table 1, the crumb hardness and gumminess of the bread containing 10% sprouted wheat sourdough were significantly (p<0.05) lower than those of the samples with 5% and 15% sourdough. Additionally, there was no significant difference in overall acceptability between the samples containing 5% and 10% sourdough. Therefore, a 10% sourdough formulation was selected as the optimal for further stages of the research. The optimal garlic extract amount in the bread formulation, based on overall acceptability scores, was selected as 7.5% as a substitute for water in the sourdough formulation.

Table 1. Textural and sensorial properties of the produced breads containing different amounts of sprouted wheat sourdough. The different letters in each column show significant difference at p<0.05.

1			
Bread sample	Overall acceptability	Gumminess	Crumb hardness (N)
5% Sourdough	2.92 ± 0.16 a	1934.20 ± 77.35 a	a 0.37 ±4.00
10% Sourdough	2.48 ± 0.43 a,b	$842.59 \pm 128.53 \text{ b}$	b 0.33 ±1.93
15% Sourdough	$2.16\pm0.29~b$	1474.60 ± 288.50 a,b	0.19 a ±3.43

3.2. Textural characteristics of the produced bread

As shown in Table 2, the addition of yogurt vinegar significantly (p < 0.05)and increased the crumb hardness while also volume. increasing its specific Additionally, sprouting significantly reduced both crumb hardness and specific volume. The inclusion of sourdough did not result in a significant reduction in crumb hardness, but it significantly increased the specific volume. Moreover, adding garlic extract to the bread significantly increased both crumb hardness and specific volume, but incorporating garlic extract into the sourdough formulation significantly reduced crumb hardness and specific volume of the produced bread.

Table 2. Overall acceptability, pH, specific volume and crumb hardness of the produced bread samples including control bread (C), bread containing sprouted wheat (SW), fermented sprouted wheat (FS), garlic extract (E), calcium propionate (CP), yoghurt and vinegar (YV) and fermented sprouted wheat containing garlic extract (FS+E). The different letters in each column indicate significant difference at p<0.05.

Bread sample	Overall	pH	Specific volume	Crumb hardness
1	acceptability	1	(cm^3/g)	(N)
С	4.40 ± 0.21 a	6.15 ± 0.02 a	$2.42 \pm 0.02 \text{ d}$	$3.83 \pm 0.02 \text{ d}$
SW	$3.68\pm0.34~b$	6.13 ± 0.01 a	$1.72\pm0.06~e$	$2.92\pm0.04~e$
FS	4.20 ± 0.17 a	$5.93\pm0.03~\mathrm{c}$	3.56 ±0.03 b	$3.55\pm0.20~d$
E	4.22 ± 0.43 a	$6.04\pm0.01~b$	2.94 ±0.09 c	$6.45\pm0.12~b$
СР	4.14 ± 0.21 a	6.13 ± 0.02 a	$2.52\pm0.02~d$	$4.70\pm0.16\ c$
YV	0.21 ±4.14 a	6.00 + 0.00 b	$3.86 \pm 0.16 \text{ a}$	$7.63\pm0.16\ c$
FS+E	$0.24\pm3.52\ b$	$5.94 \pm 0.01 \ c$	$1.93 \pm 0.03 \ e$	$2.94\pm0.24\;e$

As observed in the study by Nionelli et al. (2018), the specific volumes of the control bread, the bread with hop extract, and the sourdough bread with hop extract were reported as 2.71, 2.69, and 2.99 cm³/g, respectively, with the sourdough bread containing hop extract showing а significantly higher specific volume compared to the control sample. Additionally, the crumb hardness of these samples was reported as 2819, 2826, and 2706 g, respectively, indicating that the sourdough bread with hop extract had significantly lower crumb hardness compared to the control. However, the use of hop extract alone increased crumb hardness and reduced the specific volume of the produced bread [19].

In the research by Irakli et al. (2019), the specific volumes of the control bread, the sourdough with hop extract, and the sourdough with hop extract and rice bran were reported as 3.8, 3.1, and 3.4 cm³/g, respectively, with significant differences among them [24]. Generally, in samples containing sprouted substrate, the intensification of enzymatic activity and the breakdown of gluten and starch, which ultimately leads to less gas retention, result in a reduced specific volume. In the sample with garlic extract, a higher density mixture likely formed more stable bubbles, which, despite the increased volume, resulted in a firmer texture. The type and nature of the starter culture used in sourdough and the pH of the sourdough have also different effects on the bread's characteristics. During fermentation, metabolites such as organic acids, exopolysaccharides, and enzymes are produced, positively affecting bread texture [3]. Overall, the sample with sprouted wheat, which had less crumb hardness and specific volume, lower was not a considered suitable in terms of textural characteristics. However, the sourdough sample with less crumb hardness and higher specific volume was considered more desirable texturally.

3.3. Overall acceptability of the produced bread

The lowest overall acceptability was observed in the sample containing sprouted wheat, which showed a significant difference (p<0.05) compared to the control sample. The best sample in terms of chewability and mouth-feel was the one containing garlic extract, which did not differ significantly from the control. The bread samples did not show significant differences in taste and color, although the sample containing yogurt and vinegar differed in aroma from the others. Moreover, fermentation did not significantly the impact overall acceptability of the produced bread, but adding garlic extract to the sourdough formulation decreased the overall product acceptability.

Nionelli et al. (2018) reported that the bread samples containing hop extract had a bitter and herbal taste compared to the control. However, these samples were still considered acceptable by evaluators [19]. According to Irakli et al. (2019), the sourdough bread with hop extract did not differ significantly from the control bread. Additionally, the sourdough bread with hop

extract and rice bran had the highest overall acceptability among the tested samples and did not show a significant difference from the control bread. The acceptability scores for the control bread, the sourdough bread with hop extract, and the sourdough bread with hop extract and rice bran were reported as 3.8, 3.3, and 4.2, respectively [24]. Flavor, taste, and aroma are crucial factors in bread acceptance. It has been generally established that metabolites produced during fermentation can positively affect these factors and increase acceptability. The use of plant extracts has also a significant impact on each of these factors [3].

3.4. Inhibition of surface fungal growth

As shown in Figure 1, the highest antifungal effect was observed in the samples containing yogurt and vinegar, as well as in the sample with garlic extract alone. However, there was no significant difference (p<0.05) in the inhibitory effect of these samples compared to the bread containing calcium propionate and the sprouted wheat sourdough with garlic extract. Additionally, the greatest fungal growth among the produced breads was observed in the control and the sprouted wheat sample. According to the results, both fermentation and garlic extract reduced fungal growth, with garlic extract showing a more pronounced inhibitory effect compared to fermentation alone. Furthermore, the sprouted wheat sourdough with garlic extract provided a suitable antifungal effect compared to calcium propionate. Therefore, it can be used as a biological preservative in wheat bread processing.



Fig. 1. A: surface expansion of *A. flavus* after five days of storage on the produced bread samples. B: inhibition percentage of *A. flavus* growth on the produced bread samples including control wheat bread (C), bread containing sprouted wheat (SW), fermented sprouted wheat (FS), garlic extract (E), calcium propionate (CP), yoghurt and vinegar (YV) and fermented sprouted wheat containing garlic extract (FS+E). The different letters indicate significant difference at p<0.05.

Nionelli et al. (2018) observed that wheat bread containing hop extract weakly inhibited fungal growth in the produced bread. However, the addition of hop extract delayed the growth of *Penicillium roqueforti* for at least 14 days, which was similar to bread with 0.3% calcium propionate. They attributed the antifungal effect to the presence of weak acids in the hop extract. Nonetheless, the use of sourdough containing hop extract was more effective, primarily due to the acidification by lactic acid bacteria and the synthesis of a wide range of low molecular weight compounds such as antifungal peptides [19]. Jin et al. (2021) also noted that lactic acid bacteria could produce various organic acids with antifungal properties, although the mechanisms of some are not yet fully understood. Studies have shown that acetic acid, one of the most effective organic acids, might have antifungal effects due to its low pKa [28]. Given the antimicrobial properties of the intrinsic microorganisms in sourdough [29-32], using sourdough as a biological preservative in bakery industries could be a suitable alternative to conventional synthetic preservatives. Moreover, combining antifungal plant extracts with sourdough can significantly enhance mold control in sourdough breads while modulating the sour taste [33-35].

4- Conclusion

The use of sprouted grain sourdough in wheat bread processing is of technological significance. These ingredients not only affect the textural and sensory properties of the product but can also serve as natural additives, improving the nutritional properties and shelf-life of the bread. Additionally, enzymatic activity during sprouting facilitates the breakdown of more complex substrates into simpler compounds, which can then be utilized by microorganisms during fermentation. Therefore, sprouted grains are suitable substrates for sourdough fermentation. In this study, fermented sprouted wheat with garlic extract positively impacted the texture, overall acceptability, and mold inhibition of the produced wheat bread, with the least crumb hardness observed in the bread containing fermented sprouted wheat with garlic extract. This sourdough also effectively prevented mold growth and showed no significant difference in antifungal activity compared to calcium propionate. Based on these results, using garlic extract alone or sprouted wheat alone did not significantly impact the texture of the produced samples. However, the combined use of both in sourdough

formulation proved to be a simple and costeffective method to improve the quality attributes of the wheat bread.

5-References

- Debonne, E., Vermeulen, A., Van Bockstaele, F., Soljic, I., Eeckhout, M., & Devlieghere, F. (2019). Growth/no-growth models of in-vitro growth of *Penicillium paneum* as a function of thyme essential oil, pH, aw, temperature. *Food Microbiology*, 83, 9-17.
- [2] Ryan, L. A. M., Dal Bello, F., & Arendt, E. K. (2008). The use of sourdough fermented by antifungal LAB to reduce the amount of calcium propionate in bread. *International Journal of Food Microbiology*, 125(3), 274-278.
- [3] Sadeghi, A., Ebrahimi, M., Hajinia, F., Kharazmi, M. S., & Jafari, S. M. (2023). FoodOmics as a promising strategy to study the effects of sourdough on human health and nutrition, as well as product quality and safety; back to the future. *Trends in Food Science & Technology*. 136, 24-47.
- [4] Garcia, M. V., Bernardi, A. O., & Copetti, M. V. (2019). The fungal problem in bread production: Insights of causes, consequences, and control methods. *Current Opinion in Food Science*, 29, 1-6.
- [5] Russo, P., Fares, C., Longo, A., Spano, G., & Capozzi, V. (2017). *Lactobacillus plantarum* with broad antifungal activity as a protective starter culture for bread production. *Foods*, 6(12), 110.
- [6] Ávila Sosa Sánchez, R., Portillo-Ruiz, M. C., Viramontes-Ramos, S., Muñoz-Castellanos, L. N., & Nevárez-Moorillón, G. V. (2015). Effect of Mexican oregano (*Lippiaberlandieri* schauer) essential oil fractions on the growth of *Aspergillus* spp. in a bread model system. *Journal of Food Processing and Preservation*, 39(6), 776-783.
- [7] Hajinia, F., Sadeghi, A., & Sadeghi Mahoonak,
 A. (2021). The use of antifungal oat-sourdough lactic acid bacteria to improve safety and technological functionalities of the

supplemented wheat bread. Journal of Food Safety, 41(1), e12873.

- [8] Purabdolah, H., Sadeghi, A., Ebrahimi, M., Kashaninejad, M., Shahiri Tabarestani, H., & Mohamadzadeh, J. (2020). Techno-functional properties of the selected antifungal predominant LAB isolated from fermented acorn (*Quercus persica*). Journal of Food Measurement and Characterization, 14, 1754-1764.
- [9] Pahlavani, M., Sadeghi, A., Ebrahimi, M., Kashaninejad, M., & Moayedi, A. (2024). Application of the selected yeast isolate in type IV sourdough to produce enriched clean-label wheat bread supplemented with fermented sprouted barley. *Journal of Agriculture and Food Research*, 15, 101010.
- [10] Kia, P. S., Sadeghi, A., Kashaninejad, M., Zarali, M., & Khomeiri, M. (2024). Application of controlled fermented amaranth supplemented with purslane (*Portulaca oleracea*) powder to improve technological functionalities of wheat bread. Applied Food Research, 4(1), 100395.
- [11] Ziaee rizi, A., Sadeghi, A., Jafari, S. M., Feizi, H., & Purabdolah, H. (2024). Controlled fermented sprouted mung bean containing ginger extract as a novel bakery biopreservative for clean-label enriched wheat bread. *Journal of Agriculture and Food Research*, 16, 101218.
- [12] Zarali, M., Sadeghi, A., Ebrahimi, M., Jafari, S. M., & Mahoonak, A. S. (2024). Technonutritional capabilities of sprouted clover seeds sourdough as a potent bio-preservative against sorbate-resistant fungus in fortified clean-label wheat bread. *Journal of Food Measurement* and Characterization, 1-13.
- [13] Aryashad, M., Sadeghi, A., Nouri, M., Ebrahimi, M., Kashaninejad, M., & Aalami, M. (2023). Use of fermented sprouted mung bean (*Vigna radiata*) containing protective starter culture LAB to produce clean-label fortified wheat bread. *International Journal of Food Science & Technology*, 58(6), 3310-3320.
- [14] Rouhi, E., Sadeghi, A., Jafari, S. M., Abdolhoseini, M., & Assadpour, E. (2023).Effect of the controlled fermented quinoa

containing protective starter culture on technological characteristics of wheat bread supplemented with red lentil. *Journal of Food Science and Technology*, 60(8), 2193-2203.

- [15] Ebrahimi, M., Noori, S. M. A., Sadeghi, A., emir Coban, O., Zanganeh, J., Ghodsmofidi, S. M., ... & Raeisi, M. (2022). Application of cereal-bran sourdoughs to enhance technological functionality of white wheat bread supplemented with pumpkin (*Cucurbita pepo*) puree. *LWT*, *158*, 113079.
- [16] Ebrahimi, M., Sadeghi, A., Sarani, A., & Purabdolah, H. (2021). Enhancement of technological functionality of white wheat bread using wheat germ sourdough along with dehydrated spinach puree. *Journal of Agricultural Science and Technology*, 23(4), 839-851.
- [17] Sadeghi, A., Ebrahimi, M., Raeisi, M., & Ghods Mofidi, S. M. (2019). Improving the antioxidant capacity of bread rolls by controlled fermentation of rice bran and addition of pumpkin (*Cucurbita pepo*) puree. *Journal of Food Measurement and Characterization*, 13, 2837-2845.
- [18] Sadeghi, A., Ebrahimi, M., Mortazavi, S. A., & Abedfar, A. (2019). Application of the selected antifungal LAB isolate as a protective starter culture in pan whole-wheat sourdough bread. *Food Control*, 95, 298-307.
- [19] Nionelli, L., Pontonio, E., Gobbetti, M., & Rizzello, C. G. (2018). Use of hop extract as antifungal ingredient for bread making and selection of autochthonous resistant starters for sourdough fermentation. *International Journal* of Food Microbiology, 266, 173-182.
- [20] Hu, F., Tu, X. F., Thakur, K., Hu, F., Li, X. L., Zhang, Y. S., ... & Wei, Z. J. (2019). Comparison of antifungal activity of essential oils from different plants against three fungi. *Food and Chemical Toxicology*, 134, 110821.
- [21] AACC International. (2010). Approved methods of the American association of cereal chemists.
- [22] Montemurro, M., Pontonio, E., Gobbetti, M., & Rizzello, C. G. (2019). Investigation of the nutritional, functional and technological effects

- of the sourdough fermentation of sprouted flours. *International Journal of Food Microbiology*, 302, 47-58
- [23] Cereal and cereal products, flat bread; Sangak (2000). Institute of standards and industrial research of Iran, ISIRI 6943.
- [24] Irakli, M., Mygdalia, A., Chatzopoulou, P., & Katsantonis, D. (2019). Impact of the combination of sourdough fermentation and hop extract addition on baking properties, antioxidant capacity and phenolics bioaccessibility of rice bran-enhanced bread. *Food Chemistry*, 285, 231-239.
- [25] Katina, K., Heiniö, R. L., Autio, K., & Poutanen, K. (2006). Optimization of sourdough process for improved sensory profile and texture of wheat bread. *LWT*, 39(10), 1189-1202.
- [26] Gerez, C. L., Torino, M. I., Rollán, G., & de Valdez, G. F. (2009). Prevention of bread mould spoilage by using lactic acid bacteria with antifungal properties. *Food Control*, 20(2), 144-148.
- [27] Rizzello, C. G., Nionelli, L., Coda, R., Di Cagno, R., & Gobbetti, M. (2010). Use of sourdough fermented wheat germ for enhancing the nutritional, texture and sensory characteristics of the white bread. *European Food Research and Technology*, 230, 645-654.
- [28] Jin, J., Nguyen, T. T. H., Humayun, S., Park, S., Oh, H., Lim, S., ... & Kim, D. (2021). Characteristics of sourdough bread fermented with Pediococcus pentosaceus and Saccharomyces cerevisiae and its biopreservative effect against Aspergillus flavus. Food Chemistry, 345, 128787.
- [29] Kia Daliri, F., Sadeghi, A., Khomeiri, M., Kashaninejad, M., & Aalami, M. (2016). Evaluating the antimicrobial properties of *Lactobacillus brevis* isolated from whole barley sourdough. *Journal of Food Science and Technology (Iran)*, 15(75), 247-257.
- [30] Purabdolah, H., Sadeghi, A., Ebrahimi, M., Kashaninejad, M., & Mohamadzadeh, J.
 (2022). Evaluation of probiotic and antifungal properties of the predominant LAB isolated from fermented acorn (*Quercus persica*).

Journal of Food Science and Technology (Iran), 19(124), 171-183.

- [31] Zarali, M., Sadeghi, A., Jafari, S. M., Sadeghi Mahoonak, A., & Ebrahimi, M. (2022).
 Evaluation of antimicrobial and probiotic properties of the predominant LAB isolated from fermented germinated clover seed. *Journal of Food Science and Technology* (*Iran*), 19(123), 299-315.
- [32] Kia, S., Sadeghi, A., Kashaninejad, M., Khomeiri, M., & Zarali, M. (2023). Evaluation of probiotic properties of *Lactobacillus brevis* as the predominant LAB isolated from fermented amaranth. *Journal of Food Science and Technology (Iran)*, 19(132), 65-76.
- [33] Ebrahimi, M., Sadeghi, A., & Mortazavi, S. A. (2020). The use of cyclic dipeptide producing LAB with potent anti-aflatoxigenic capability to improve techno-functional properties of clean-label bread. Annals of microbiology, 70, 1-12.
- [34] Ziaee rizi, A., Sadeghi, A., Feizi, H., Jafari, S. M., & Purabdolah, H. (2024). Evaluation of textural, sensorial and shelf-life characteristics of bread produced with mung bean sourdough and saffron petal extract. *Journal of Food Science and Technology (Iran)*, 21(148), 141-153.
- [35] Sadeghi, A., Ebrahimi, M., Assadpour, E., & Jafari, S. M. (2023). Recent advances in probiotic breads; a market trend in the functional bakery products. *Critical Reviews in Food Science and Nutrition*, 1-12.

مجله علوم و صنایع غذایی ایران

سایت مجله: <u>www.fsct.modares.ac.ir</u>

مقاله علم<u>ى پژو</u>هشى

تاثیر استفاده از خمیرترش سنتی بر ویژگیهای کیفی نان حجیم حاوی گندم جوانهزده و عصاره سیر

سارا صفرى، عليرضا صادقي*، على مؤيدى، حسين پورعبدالله، عليرضا صادقي ماهونك

نابع طبیعی گرگان، گرگان، ایران	دانشگاه علوم کشاورزی و م	مهندسی صنایع غذایی،	گروه علوم و
--------------------------------	--------------------------	---------------------	-------------

اطلاعات مقاله	چکیدہ
تاریخ های مقاله :	استفاده از عصارههای گیاهی در فرمولاسیون خمیرترش میتواند سبب بهبود ویژگیهای حسی و
تاریخ دریافت: ۱٤،۴۲/٤/۲۷	زمان ماندگاری نان تولیدی گردد. در پژوهش حاضر، تاثیر استفاده توام از گندم جوانهزده تخمیر
تاریخ پذیرش: ۱٤۰۳/۸/۲۰	شده و عصاره سیر در فرمولاسیون خمیرترش سنتی حاوی ماست و سرکه بر ویژگیهای کیفی نان
	گندم حجیم مورد بررسی قرار گرفت. بر اساس نتایج بافتسنجی از بین نمونههای تولیدی، کمترین
کلمات کلیدی:	سفتی بافت در نان حاوی گندم جوانهزده تخمیر شده و عصاره سیر (۲/۹٤ نیوتن) مشاهده شد که
عصارہ سیر،	به شکل معنیداری (p<۰/۰۵) از سفتی بافت نمونه شاهد کمتر بود. استفاده توام از تخمیر و
گندم جوانەزدە تخمير شدە،	جوانهزنی نیز سبب افزایش حجم مخصوص و کاهش چسبندگی بافت نان تولیدی گردید. علاوه
خمیرترش سنتی، سفتے یافت،	بر این، اگر چه کمترین توسعه سطحی قارچ Aspergillus flavus در نانهای حاوی ماست و سرکه
ى . كېكىزدگى.	و همچنین عصاره سیر به تنهایی مشاهده شد اما تفاوت معنیداری بین میزان توسعه سطحی قارچ
	در نمونه حاوی گندم جوانهزده تخمیر شده و عصاره سیر با نان حاوی پروپیونات کلسیم وجود
DOI:10.22034/FSCT.22.159.133.	نداشت. همچنین نمونه حاوی گندم جوانهزده تخمیر شده با نمونه شاهد به لحاظ پذیرش کلی
* مسئول مكاتبات:	تفاوت معنیداری نشان نداد. بر این اساس، خمیرترش گندم جوانهزده حاوی عصاره سیر به عنوان
sadeghi.gau@gmail.com	یک نگهدارنده زیستی میتواند جهت بهبود ویژگیهای کیفی و زمان ماندگاری نان گندم مورد
	استفاده قرار گیرد.