



Scientific Research

Effect of Bromhexine Supplementation on Reproductive Performance and its Implications for Meat and Milk Production in Damascus Goat Bucks during the Non-Breeding Season

Layla Abd-Alsattar Sadiq Laylani¹, Ansam Hussein Ali², Yahya Waheeb Ibrahim

1. Community Health Department, Technical Medical Institute Kirkuk, Northern Technical University

2. Northern Technical University, Al-dor Technical Institute, Department of pharmacy techniques

3. Department of Biotechnology, College of Applied Sciences, University of Samarra

ARTICLE INFO	ABSTRACT
<p>Article History:</p> <p>Received: 2025/12/28 Accepted: 2026/2/6</p>	<p>The dynamics of reproductive performances of Damascus goats through seasonal changes are a major challenge to the production of milk and meat during year-round, which influences the sustainability of all the supply chains. Mucolytic agent Bromhexine has demonstrated a possible role in enhancing sperm quality by decreasing and viscosity therefore indirectly increasing the breeding efficiency and subsequently production of animal-derived foods. The present study examined the impact of bromhexine supplementation on the semen quality parameters in Damascus goat bucks during non-breeding season and implications of such research on reproductive management and subsequent impacts on milk and meat production. A total of twelve sexually fit Damascus bucks (2-3 years old, 45-55 kg) were randomly divided in a treatment group with oral bromhexine (1 mg/kg day) or a control group with placebo after 60 days during the non-breeding season. Sperm was pooled and assessed on volume, liquefaction period, viscosity, sperm density, motility, viability as well as morphology. The serum reproductive hormones and semen plasma biochemical constituents were also evaluated. Bromhexine massively lowered the liquefaction time (8.2±1.1 min vs. 15.7±1.8 min, p<0.001) and seminal plasma viscosity (2.8±0.3 cP vs. 4.6±0.5 cP, P<0.01). There was an improvement in progressive sperm motility (66.3±3.4% vs. 52.8±4.2%, p<0.05). There were no major differences in sperm concentration, viability, morphology, hormonal profiles or biochemistry of the seminal plasma. Bromhexine enhances some of the key semen physical characteristics, and this might increase the breeding efficiency in the non-breeding season. This can be used to improve year-round breeding that will result in stabilized production of milk and meat. The fact that there are no hormonal changes is an indication that bromhexine has no effect on the normal physiology of food producing animals, hence it is a possible alternative that can be incorporated into sustainable animal husbandry practices in an attempt to streamline the food supply chain resilience.</p>
<p>Keywords:</p> <p>Bromhexine, Damascus goat, semen quality, meat production, milk supply, food security, animal husbandry</p>	
<p>DOI: 10.48311/fsct.2026.118564.83009.</p> <p>*Corresponding Author E-Mail: Ansam.ha@ntu.edu.iq</p>	

1. Introduction

The stability and sustainability of animal-derived food supply chains represent a cornerstone of global food security and economic viability within the agricultural sector. Small ruminants, particularly goats, play an indispensable role in this context, serving as a primary source of high-quality protein, essential fats, and micronutrients for a significant portion of the world's population, especially in arid and semi-arid regions [1]. One of the most praised breeds, the Damascus goat (*Capra hircus*), also called the Shami goat, is characterized by a high dairy production level, the high quality of the meat, and the ability to survive under the most harsh environmental conditions [2]. The breed is one of the significant agricultural resources of the Middle East and the Mediterranean basins and the main source of livelihood of most of the local economies and subsistence farming systems. The productivity of this valuable genetic resource is however restricted severely by a basic biological constraint, that of extreme seasonal reproductive patterns [3]. Similar to most photoperiod sensitive species living in subtropical and temperate areas, Damascus goats are highly seasonal in reproductive actions, with the sophisticated interaction between photoperiod (daylight length) and the neuroendocrine axis [4]. This seasonality is most highly manifested in the long-day photoperiods of spring and early summer, the time unanimously accepted as the non-breeding season of this breed in the Northern Hemisphere [5]. This stage is marked by a major reduction in reproductive performance, which consists of a decreased libido, the decrease in testicular volume, the decrease in the amount of testosterone released and a drastic decline in the parameters of semen quality [6]. The most seminal features that have been most negatively affected are the ejaculate volume, sperm concentration, progressive motility and most importantly the physical properties of the semen itself such as higher viscosity and extended liquefaction.

time [7]. These are changes that are not just theoretical findings in books, but which are also directly translated into practical barriers to breeding programmes. It is the high viscosity and slow liquefaction, which physically encloses spermatozoa in a mucoid network and limits their liberated release and subsequent motility, although the sperm cells may still have inherent functional competence [8]. This means that the fertility of ejaculates gathered during non-breeding season is plummeted and thus, fewer conception rates occur through natural copulation as well as artificial insemination (AI) [9]. In the viewpoint of food systems and industrial farming, such reproductive seasonality is a daunting challenge to the reliability and efficiency of goat meat (chevon) and goat milk production. Its immediate result is a very seasonalized and peak supply of children and milk, which results in a cycle of feast and famine in the market [10]. This imbalance interferes with the level flow of raw materials to the processing facilities, inconveniences stock control to the retailers, and causes fluctuations in prices which are no benefit to the producer or the consumer [11]. In the case of dairy operations, the inability to have year-long breeding patterns, causes seasonal milk production curves, which forces processors to respond to a changing milk intake, posing inefficiencies in the capacity utilization [12]. With a narrow band of concentration in kidding in a meat production system the animals are over-slaughtered at certain intervals causing the prices in the market to fall and a short time later there are periods of scarcity [13]. Mitigating seasonal infertility is thus not only an aim of reproductive physiology, but a major necessity in enhancement of the resilience, profitability and sustainability of the overall small ruminant food sector.

The best-known analogous approaches to seasonal reproductive depression have been endocrine axis manipulation. They involve exogenous hormone administration (e.g. melatonin, gonadotropin-releasing hormone (GnRH) analogs, and prostaglandins) or the use of exogenous hormone administration (e.g. gonadotropin-releasing hormone (GnRH) analogs and prostaglandins).

regulated light photoperiod with lightproofing [14]. These approaches may be good, but tend to have a lot of practical and financial constraints. Hormonal treatment is also associated with residues accumulating in the animal products and this is a major problem in a fast maturing consumer-focused environment and withdrawal may be too long making it difficult to manage schedules [15]. Moreover, there may be regulatory challenges and consumer distrust in the use of hormonal interventions, which may restrict access to the market [16]. The manipulation of photoperiod, in its turn, requires a significant amount of capital investment in specialized infrastructure and raises energy costs of operations, which is not in the capability of a number of small to medium-sized producers [17]. This generates an interesting interest in other, non-hormonal, and cost-effective interventions, which can be used to increase reproductive efficacy in the non-breeding season and prevent these side effects. It is against this background that the possibilities of bromhexine hydrochloride are worthy of investigation. Bromhexine (2-amino-3,5-dibromo-N-cyclohexyl-N-methylbenzylamine) is a long-established mucolytic agent, initially developed and used with great success in the treatment of respiratory disorders, which are characterised by excessive and viscous mucus [18]. The main mechanism of action of it is the depolymerization of the mucopolysaccharide fibres, namely disulfide bonds, lowering of the molecular weight of complex glycoproteins, leading to the decrease of the viscosity of respiratory secretions and clearance of them [19]. The bromhexine chemical structure is able to react with glycoprotein complexes not only in the respiratory tract but also in other biological fluids [20]. Since seminal plasma of goats and other mammals has large amounts of mucoproteins, glycosaminoglycans and semenogelin proteins that influence the development of the coagulum post-ejaculation and total semen viscosity [21], it is biologically possible that bromhexine will have the same

mucolytic effect.

within the seminal plasma. Bromhexine has the potential to depolymerize these mucoid constituents and therefore hasten the liquefaction of semen, decrease residual viscosity and literally remove spermatozoa, thus raising their observable motility without necessarily compromising their inherent biological quality or the hormonal milieu per se [22]. The suggested peripheral mechanism of action, specifically, is especially appealing to the food industry perspective given that it would avoid the necessity of direct hormonal action, in turn, eliminating the apprehension of hormonal residues in meat and milk and conforming to consumer demands of the kind of cleaner production. Other species of evidence give supportive but still preliminary justification in this approach. Ram studies have also shown that the administration of bromhexine is capable of reducing semen coagulation time and enhancing progressive motility, and this effect is more pronounced in the non-breeding season [23]. Bromhexine treatment in bulls was linked to a better semen flowability and processing properties, which are vital in AI and semen cryopreservation [24]. Likewise, a study conducted on roosters has demonstrated that bromhexine supplementation has a potential of reducing semen quality deterioration caused by heat stress such as viscosity decrease and motility enhancement [25].

These interspecies results indicate that fundamental mechanism is preserved which would be useful to exploit in Damascus goats. Nevertheless, there is a research gap that is critical. Although the bromhexine influence on semen physical properties is encouraging, there is no profound study in the peculiar environment of Damascus goat bucks in the non-breeding period, and with the strict regard to the repercussions of meat and milk production systems. It has yet to be clearly established how the positive changes in the parameters of semen can be translated into an effective instrument to stabilize the production cycles on an annual basis. Moreover, one should ensure that the effect of bromhexine is peripheral and does not disrupt the hypothalamic-pituitary-gonadal.

This would support the safety profile of (HPG) axis to be used in food-producing animals. It would also be highly desirable to confirm that it has no effects on important seminal plasma biomarkers of accessory sex gland activity (e.g. fructose of seminal vesicles, citric acid of the prostate) in order to conclude that its effects are purely physical rather than secretory [26]. Hence, the research was aimed at filling this gap by providing an intensive assessment of the impacts of oral bromhexine supplement on a holistic set of semen quality indices in Damascus goat bucks in the non-breeding season. We used the best analytical methods such as computer-assisted sperm analysis (CASA) to acquire accurate motility cycles and standardized rheological methods of viscosity to gain an in-depth analysis of the treatment effects [28,29]. Simultaneously, we were observing systemic levels of reproductive hormones as well as seminal plasma biochemistry to define the mechanism of action. We predict that the activity of bromhexine will be a mucolytic agent in the seminal plasma, which reduces the liquefaction time and viscosity with considerable influence on the sperm motility parameters due to the improved physical constraints, and bromhexine will not cause any significant changes in the process of spermatogenesis, sperm viability, the endocrine activity, or the biochemical composition of the seminal plasma. It is expected that the results of this study would serve as a scientifically valid, practical, and safe approach to improving the reproductive efficiency of one of the primary dairy and meat breeds, which would directly lead to the stabilization of animal protein sources, decrease the production variability in the seasons, and encourage a higher level of sustainability in the goat breeding industry.

2. Materials and Methods

2.1 Ethical Considerations

This study was conducted in accordance with institutional policies for the care and use of animals, which are consistent with the principles outlined in the Guide for the Care and Use of Agricultural Animals in Research and Teaching; All the protocols of experiments were reviewed and approved by Institutional Animal Care and Use Committee (Approval No. IACUC-2024-037). Animals were humanely handled during the experimental time and procedures were designed to minimize stress and discomfort.

2.2 Experimental Animals and Management

Twelve Damascus bucks of sexually mature age, 2-3 years of age body weights of 45-55 kg were chosen from a commercial goat farm at position 35deg N latitude. Bucks were selected because of prior examination for breeding soundness based upon evidence of normal reproductive ability and semen quality during the breeding season. Animals were kept in individual pens (2.5 x 3 m) with natural photoperiod, having free access to fresh water. The diet was consisted of concentrate mixture designed to satisfy nutritional requirements of breeding males (16% crude protein, 2.8 Mcal/kg metabolizable energy), alfalfa hay and seasonal green fodder. All animals were vaccinated following regional protocols and also dewormed with ivermectin two weeks before the commencement of the experiment. Bucks had no clinical diseases during the experimental time, as could be verified through weekly veterinary examinations.

2.3 Experimental Design and Treatment

The experiment was conducted during the non-breeding season (April through June), corresponding to the period of long-day photoperiods (average 14.2 hours daylight) and naturally reduced reproductive activity in this breed. Following a two-week adaptation period during which all animals were trained for semen collection, bucks were randomly allocated to two equal groups (n=6 per group) using a completely randomized design:

Treatment Group (BRX): Bucks received oral bromhexine hydrochloride (Bisolvon®, Boehringer Ingelheim, pharmaceutical grade purity >98%) at a dose of 1 mg/kg body weight daily. The drug was

dissolved in 10 ml normal saline and administered orally using a dosing syringe each morning at 0800 h before feeding. Body weights were recorded weekly and doses adjusted accordingly.

Control Group (CON): Bucks received an equivalent volume of oral placebo (normal saline, 10 ml) using the same administration protocol to control for handling effects.

The duration of treatment was 60 days, which corresponds to about two complete spermatogenic cycles in goats about 49 days of time from spermatozoa of spermatogonia to fully mature spermatozoa so that all spermatozoa tested after treatment had been exposed to bromhexine during their development and epididymal maturation.

2.4 Semen Collection and Initial Evaluation

Semen samples were collected twice a week (Tuesday and Friday mornings between 0900-1100h) using an artificial vagina with a temperature of 42-45degC using an estrous doe as teaser. Collection frequency and timing were standardized in order to minimize variation as a consequence of an occurrence of sexual rest periods. To each collection, bucks were presented with 2 false mounts followed with semen collection on the 3rd mount in order to insure optimal sexual stimulation. Immediately after collection samples were protected from light and temperature shock by allowing collection tubes to be contained in a portable water bath maintained at 37degC.

Initial macroscopic evaluation was done in less than 2 minutes of collection and included:

Ejaculate Volume: Measured straight from the graduated collection tube to the nearest 0.05 ml from the fraction (if any) of the gel.

Color and Consistency: Recorded on standardized descriptors (color: white, creamy-white, yellowish; consistency: watery, normal, thick, very thick).

pH: measured with calibrated digital pH meter (Hanna instruments, HI98103) at 37oC immediately after collection

2.5 Liquefaction Time Assessment

Liquefaction time, which is the time elapsed from ejaculation to the time when the semen began flowing freely as single droplets, was evaluated using the standardized method for tilting [31]. The sample tube was immediately tipped over at 45o every minute. Complete liquefaction was noted when semen was released in separate drops and not viscous threads. Liquefaction time was measured in minutes from ejaculation time. This parameter is important in understanding the main mechanism of action of bromhexine.

2.6 Seminal Plasma Viscosity Measurement

Seminal plasma viscosity was tested with a calibrated Ostwald viscometer kept at 37degC in a water bath [32]. After liquefaction, 2 ml of semen was centrifuged at 3000 x g for 10 minutes at room temperature to separate seminal plasma and spermatozoa. The supernatant was carefully collected without interfering with the sperm pellet. Viscosity was measured by measuring the flow time of seminal plasma and comparing it with distilled water (reference standard) in the viscometer capillary. Dynamic viscosity was determined from the formula:
$$\eta = (t \times \rho \times K)$$

where η = dynamic viscosity (centipoise, cP), t = time of flow (sec), ρ = density (assumed = 1.0 g/ml for seminal plasma), and K = constant of viscometer (calculated from calibration of viscometer with standard fluids). All measurements were done in duplicates and the mean value was determined.

2.7 Sperm Motility Analysis

Progressive sperm motility was evaluated using both subjective and objective computerized methods:

Subjective Assessment: After complete liquefaction, 10ml of semen was diluted 1:20 in pre-warmed (37degC) phosphate buffered saline (PBS, pH 7.2). A 5 ml drop was placed on a microscope slide that was previously warmed and covered with a coverslip that was 22x22 mm in size. Progressive motility was measured at 400x magnification with the help of phase-contrast microscopy (Olympus CX43) with a heated stage at 37degC. At least 200 spermatozoa per sample in five random fields was evaluated by one experienced technician who was blinded as to treatment groups. Spermatozoa were either classified as progressively motile (forward movement, regardless of speed), non-progressively motile (all other movement patterns) or immotile. Results were calculated as percentage of progressively motile spermatozoa.

Computer-Assisted Sperm Analysis (CASA): Sperm kinematic parameters were objectively analyzed as in a computer-assisted sperm analysis (CASA) system (AndroVision, Minitube GmbH, Germany) with the following parameters optimized for caprine spermatozoa: frame rate of 60 Hz, minimum 25 frames captured in each field, detection parameters of the head size of the goat spermatozoa set at 30-50 μm^2 . Samples were diluted 1:20 in a preparation buffer (pre-warmed Tris-citrate extender (pH 6.8)) and loaded into a Leja 4-chamber slide of 20mm depth. At least 7 microscopic fields and 500 spermatozoa per sample were analysed at 37degC. Parameters that were evaluated were:

Total motility, TM (%): percent motility of spermatozoa exhibiting any motility

Progressive motility (PM, %): % showing forward progression (VAP >25 $\mu\text{m}/\text{s}$ and STR >80%)

Curvilinear velocity VCL, $\mu\text{m}/\text{s}$ Total
time = distance traveled divided by time

Straight-line velocity (VSL, $\mu\text{m}/\text{s}$) Straight-line distance from start to end point divided by time

Average path velocity (VAP, $\mu\text{m}/\text{s}$): velocity along to the smoothed average path

Linearity (LIN, %): (VSL/VCL x 100)
Measuring deviation from straight line

Straightness (STR, %): (VSL/VAP)x100,
deviation from average path

Wobble (WOB, %): (VAP/VCL) x 100,
oscillation of actual path measuring

Amplitude of lateral head displacement (ALH, μm)
Magnitude of lateral head movement

Beat cross frequency (BCF, Hz) Frequency of flagellar beating

2.8 Sperm Concentration Determination

Sperm concentration was performed with a hemocytometer technique with spectrophotometric analysis confirmation:

Hemocytometer Method: Semen was diluted 1:100 with formol-saline solution (contain formol - 4% and NaCl - 0.9% for the immobilization of spermatozoa) and mixed well. After 5 minutes of equilibration time, the diluted sample was placed in both chambers of an improved Neubauer hemocytometer. Spermatozoa were enumerated in five large squares (1mm² each) in both chambers under 400x magnification. The concentration was calculated by using the standard formula and expressed as x10⁹ spermatozoa/ml. Samples with the coefficient of variation

Recounted >15% between chambers were recounted.

Spectrophotometric Verification: The sperm concentration was also estimated by using a photometer (Accucell(R), IMV Technologies, France) calibrated specially for caprine semen as per the protocol of the manufacturer. This method resulted in fast verification of hemocytometer results.

2.9 Sperm Viability Assessment

The viability of sperm was evaluated using the method of eosin-nigrosin staining [33]. A 10 ml aliquot of liquefied semen was mixed with 20 ml of eosin-nigrosin stain (1% eosin Y and 10% nigrosin in distilled water) on a pre-warmed glass slide at 37degC. A thin smear was prepared by gently mixing for 30 seconds and left to air dry at room temperature using the edge of another slide. Slides were examined within 2 hours at 1000x with oil immersion using bright-field microscopy. At least 200 spermatozoa for each sample was evaluated in random fields. Spermatozoa whose plasma membranes are intact exclude the eosin dye and appear white (live) while those with compromised membranes take up the stain and appear

pink to red (dead). Results were reported as per cent viable (unstained) spermatozoa. Hypo-osmotic Swelling

Test (HOST) 2.10

Plasma membrane functional integrity was measured by the hypo-osmotic swelling test [34]. A 100ml aliquot of liquefied semen was gently mixed with 1ml of pre-warmed (37°C) hypo-osmotic solution (100mOsm/kg, consisting of 0.735g sodium citrate dihydrate and 1.351g D-fructose dissolved in 100ml distilled water, osmolality verified using a freezing point osmometer). Samples were incubated at 37°C for 60 min. in a water bath. After incubation, a drop of 10ml was placed on a microscope slide and observed at 400x using phase-contrast microscope. At least 200 spermatozoa in random fields were considered. Spermatozoa with functional plasma membranes react to hypo-osmotic stress by the influx of water producing characteristic coiling or tail swelling (classified as g-type curling). Non-responsive spermatozoa with damaged membrane function have normal Straight tail morphology. Results were given as percentage of HOST-positive spermatozoa.

2.10 Sperms Morphological Evaluation

Sperm morphology was evaluated from eosin-nigrosin stained smears made for viability evaluation. At least 200 spermatozoa per sample were studied at 1000x magnification at oil immersion in bright-field. Morphological abnormalities were classified following standard veterinary andrological procedures in the following three main categories:

Head abnormalities: Pyriform (pear shaped), narrow at the base, small or large heads, double heads, detached heads, abnormal acrosome

Midpiece abnormalities: Bent/ folded midpiece, thickened midpiece (suggesting defects in mitochondrial sheath), proximal cytoplasmic droplets (found at junction of head and midpiece), distal cytoplasmic droplets (found along midpiece or at midpiece-principal piece junction)

Tail abnormalities Coiled or bent tail, Short tail, Detached tail, Double tail

Normal spermatozoa were determined as spermatozoa with normal shape of their oval heads, intact acrosome, without droplets in their midpiece and principal piece, and no broken-off parts. Total abnormal spermatozoa percentage was obtained by addition of all the categories of abnormality. Individual types of abnormality were separately recorded as well.

2.11 Seminal Plasma Biochemical Analysis

For biochemical analyses, semen samples of each buck were centrifuged at 3000*g for 15 min at 4°C within 30 min of collection. The seminal plasma supernatant was carefully separated without disturbing the sperm pellet in aliquots into 1.5 ml cryovials and immediately frozen at -80°C until analyzed in batches. All samples of a given parameter were run in the same assay to eliminate any inter-assay variation.

Total Protein Concentration: Determined using the Bradford protein assay [35] using reagent from Sigma-Aldrich. Bovine serum albumin was used as standard (range 0.1 - 1.0 mg/ml). Absorbance was then read at 595 nm with the help of a spectrophotometer. Results have been expressed in mg/ml.

Fructose Concentration: Determined according to the resorcinol colorimetric method [36]. Fructose, which is secreted mainly by seminal vesicles in response to androgens, is an indicator of seminal vesicle secretory function and is the major energy substrate for the spermatozoa. A normal curve was prepared with the solutions of D-fructose (0-5 mg/ml). Results were given in mg/ml seminal plasma.

Citric Acid Concentration: Measured according to enzymatic method with the help of commercial kit (Sigma-Aldrich, MAK057). The main secretion of citric acid is from the prostate gland and is used as a marker of prostatic function. Results were expressed as mg/ml seminal plasma.

2.12 Hormone Assays

Blood samples (10 ml) were collected via jugular venipuncture from all bucks at three times: immediately before the start of treatment (Day 0), at the mid-point of treatment (Day 30) and at the conclusion of the experiment (Day 60). Samples were obtained in plain vacuum tubes between 0800-0900 h in order to minimize circadian variation in hormone concentrations. Blood was allowed to clot at room temperature for 30 minutes, centrifuged at 3000

x g for 15 minutes. Serum was separated and aliquoted and stored at -20°C until analyzed. All samples were analysed in duplicates in the same assay batch.

Testosterone Concentration: Measured by enzyme-linked immunosorbent assay (ELISA) with commercial kits DRG International Inc USA, Catalog No. EIA-1559 validated in previously caprine serum. The assay is performed on the principle of competitive binding of testosterone in samples and horseradish peroxidase

conjugated testosterone for a fixed number of antibody binding sites. Intra-assay coefficient of variation was 4.2%, inter-assay coefficient of variation was 7.8% and analytical sensitivity was 0.083 ng/ml. Results were expressed as ng/ml.

Follicle-Stimulating Hormone (FSH): Determined using a competitive enzyme immunoassay (ELI) (ELS) kit (MyBioSource Inc. USA, Catalog no. MBS269090) validated for ruminant species. Intra-assay CV was 5.1%, inter-assay CV was 9.2% and sensitivity was 0.5mIU/ml. Amount was expressed in mIU/ml.

Luteinizing Hormone (LH): Measured by using Enzyme Immuno-Assay (ELISA) using commercial kits, Cusabio Technology LLC USA, Catalog No. CSB-E06869Cp, using caprine serum as validated following manufacture's protocols. Intra-assay CV was 4.8%, inter-assay CV was 8.5% and sensitivity was calculated as 0.39 mIU/ml. Results were expressed as mIU/ml.

2.13 Statistical Analysis

All data were first checked for outliers with Grubbs' test ($\alpha=0.05$). Data were then tested for normality using the Shapiro-Wilk statistical test and homogeneity of variance using Levene's test. Data satisfying parametric assumptions were analyzed by repeated measures analysis of variance (ANOVA) with treatment as the between-subject factor and collection time (weeks) as the within-subject factor, and the mixed procedure in SAS (SAS Institute Inc. version 9.4, Cary, NC, USA) The statistical model was treatment, time and treatment x time interaction in the fixed effects and individual animal in the random effect.

For the semen parameters collected more than once per week, data were averaged per week for each animal before statistical analysis to prevent pseudoreplication. Where significant main effects or interactions were found ($P<0.05$) pairwise comparisons of least square means were calculated using the Tukey's honestly significant difference (HSD) adjustment for multiple comparisons.

Data which did not conform to parametric assumptions after attempts to transform them (logarithmic, square root, or arcsine transformations)

were analyzed using non-parametric alternatives. The Mann-Whitney U test was applied to compare two independent groups and the Kruskal-Wallis test and Dunn's post-hoc test were applied to compare more than two groups.

Hormone concentration data were analyzed with the repeated measures of analysis of covariance (ANOVA) with 3 time points (Day 0, 30, 60). When the violation of sphericity assumption was observed (Mauchly's test, $P<0.05$), Greenhouse-Geisser correction was used.

Pearson correlation coefficients were determined to analyze relationships between the parameters of semen quality. Viscosity, liquefaction time and motility parameters were analyzed in particular relationship between them. Correlation matrices were developed and displayed with GraphPad Prism (version 9.0, GraphPad Software, San Diego, CA, USA).

All statistical analyses were two-tailed and statistical significance was defined as $P<0.05$. Trends were discussed at $0.05<P<0.10$. Results are presented in terms of least squares mean±standard error of the mean (SEM), unless otherwise stated.

Sample size determination was based on previous studies looking at the effects of bromhexine on ram semen

quality [22], with power analysis showing that $n=6$ per group would give 80% power of detecting a difference of 15% in progressive motility with $\alpha=0.05$

3.3. Results

3.1 Macroscopic characteristics of the Semen

Macroscopic semen characteristics of Damascus bucks during the non-breeding season are given in

Table 1. Ejaculate volume did not show a significant difference between groups (1.08±0.09 ml and 0.95±0.08 ml in BRX and CON, respectively, $P=0.31$). All samples had creamy-white colouration typical for caprine semen with no differences in colour between groups. Semen pH was not statistically different (6.85±0.12 in BRX and 6.82±0.11 in CON, $P=0.86$). However, subjective assessment of consistency showed that bromhexine-treated samples were easier to handle and demonstrated improved flowability during the laboratory processing but this was not formally quantified.

Table 1. Effect of Bromhexine on Macroscopic Semen Characteristics

Parameter	Control	Bromhexine	P-value
Ejaculate volume (ml)	0.95±0.08	1.08±0.09	0.31
pH	6.82±0.11	6.85±0.12	0.86
Color	Creamy-white	Creamy-white	-
Gross motility (0-5)	3.2±0.3	3.7±0.3	0.24

Values are least squares means ± SEM

3.1 Liquefaction Time and Seminal Plasma Viscosity

Bromhexine supplementation had a dramatic effect on decreasing the time to liquefaction of the semen, compared with controls (Table 2). Mean liquefaction time in BRX group was

8.2±1.1 minutes significantly less than 15.7±1.8 minutes in the CON group ($P<0.001$). This is a 48% decrease in the liquefaction time, which suggests

great improvement of liquefaction process (Figure 1).

Seminal plasma viscosity (after complete liquefaction) was significantly lower in bromhexine treated bucks (2.8±0.3 cp vs. 4.6±0.5 cp, $p<0.01$). This 39% decrease of viscosity confirms the mucolytic action of bromhexine on seminal plasma constituents. The temporal pattern revealed that these effects were apparent following a period of about 14-21 days of treatment, and continued throughout the rest of the experimental period.

Table 2. Effect of Bromhexine on Liquefaction Time and Seminal Plasma Viscosity

Parameter	Control	Bromhexine	P-value
Liquefaction time (min)	15.7±1.8 ^a	8.2±1.1 ^b	<0.001
Seminal plasma viscosity (cP)	4.6±0.5 ^a	2.8±0.3 ^b	<0.01

Values are least squares means ± SEM. ^{a,b} Different superscripts within rows indicate significant differences ($p<0.05$).

3.2 Sperm Concentration and Total Sperm Output

Sperm concentration showed no significant difference between treatment groups (Table 3). The mean concentration in the BRX group was $3.2±0.4×10^9/ml$ compared to $2.9±0.3×10^9/ml$ in the CON group ($p=0.58$). Consequently, total sperm output per ejaculate, calculated as volume × concentration, also did not differ significantly ($3.46±0.52×10^9$ in BRX vs. $2.76±0.38×10^9$ in CON, $p=0.29$). These findings

indicate that bromhexine's effects did not extend to spermatogenic processes or testicular sperm production.

Table 3. Effect of Bromhexine on Sperm Concentration

Parameter	Control	Bromhexine	P-value
Sperm concentration ($×10^9/ml$)	2.9±0.3	3.2±0.4	0.58
Total sperm per ejaculate ($×10^9$)	2.76±0.38	3.46±0.52	0.29

Values are least squares means ± SEM

3.4 Sperm Motility Parameters

Progressive sperm motility, assessed by subjective evaluation, was significantly improved in the bromhexine treatment group ($66.3 \pm 3.4\%$ vs. $52.8 \pm 4.2\%$ in controls, $p < 0.05$) (Table 4). This 26% relative improvement represents the most functionally important finding of the study, as progressive motility is a critical predictor of fertility.

Computer-assisted sperm analysis indicated that bromhexine treatment significantly improved straight-line velocity (VSL: 52.4 ± 3.8 mm/s vs. 41.7 ± 3.2 mm/s, $p < 0.05$) which represents the improvement of linear progressive movement. Average path velocity (VAP) was shown to

similar trend (68.5 ± 4.1 mm/s vs. 59.2 ± 3.7 mm/s, $p = 0.09$). However, there was no difference in curvilinear velocity (VCL) (total distance traveled; 112.5 ± 7.2 mm/s in BRX vs. 105.8 ± 6.8 mm/s in CON, $p = 0.50$).

Linearity (LIN) and straightness (STR) indices improved in the BRX group, but only STR was significantly improved ($75.8 \pm 2.4\%$ vs. $70.2 \pm 2.8\%$, $p < 0.05$). Importantly, wobble (WOB) and beat cross frequency (BCF) reflecting intrinsic flagellar beat patterns, showed no significant group differences (WOB: $60.9 \pm 2.1\%$ vs. $59.6 \pm 2.3\%$, $P = 0.68$; BCF: 7.8 -

Being able to observe beats indicates that motor proteins rotate without bias, traveling across the flagellum freely without interruption by other stages of the flagellum. $P = 0.72$), showing that bromhexine did not affect the fundamental motility apparatus of spermatozoa.

Table 4. Effect of Bromhexine on Sperm Motility Parameters

Parameter	Control	Bromhexine	P-value
Progressive motility (%)	52.8±4.2 ^b	66.3±3.4 ^a	<0.05
Total motility (%)	64.5±4.8	74.2±4.1	0.15
VCL (µm/s)	105.8±6.8	112.5±7.2	0.50
VSL (µm/s)	41.7±3.2 ^b	52.4±3.8 ^a	<0.05
VAP (µm/s)	59.2±3.7	68.5±4.1	0.09
LIN (%)	39.4±2.6	46.6±2.9	0.07
STR (%)	70.2±2.8 ^b	75.8±2.4 ^a	<0.05
WOB (%)	59.6±2.3	60.9±2.1	0.68
ALH (µm)	3.8±0.4	3.6±0.3	0.72
BCF (Hz)	7.5±0.5	7.8±0.6	0.72

VCL: curvilinear velocity; VSL: straight-line velocity; VAP: average path velocity; LIN: linearity; STR: straightness; WOB: wobble; ALH: amplitude of lateral head displacement; BCF: beat cross frequency. Values are least squares means ± SEM. ^{a,b} Different superscripts within rows indicate significant differences (p<0.05)

3.5 Sperm Viability and Membrane Integrity

Sperm viability, assessed by eosin-nigrosin staining, did not differ significantly between groups (Table 5). The percentage of viable spermatozoa was 77.4±3.2% in the BRX group compared to 73.8±3.8% in controls

viability status.

(p=0.48). Similarly, the hypo-osmotic swelling test revealed no significant difference in plasma membrane functional integrity (71.2±3.5% HOST-positive in BRX vs. 67.5±4.1% in CON, p=0.51). These findings indicate that bromhexine treatment did not substantially affect sperm membrane integrity or

Table 5. Sperm Viability and Membrane Integrity

Parameter	Control	Bromhexine	P-value
Sperm viability (%)	73.8±3.8	77.4±3.2	0.48
HOST-positive (%)	67.5±4.1	71.2±3.5	0.51

HOST: hypo-osmotic swelling test. Values are least squares means ± SEM

3.6 Sperm Morphology

Total sperm morphological abnormalities were statistically similar between groups (Table 6). The BRX group exhibited 11.8±1.4% total abnormalities compared to 13.7±1.6% in the CON group (p=0.40). When categorized by abnormality type, no significant differences were observed for head abnormalities (3.2±0.6% vs. 3.8±0.7%, p=0.54), midpiece

abnormalities (4.5±0.8% vs. 5.2±0.9%, p=0.58), or tail abnormalities (4.1±0.7% vs. 4.7±0.8%, p=0.60).

Cytoplasmic droplet retention, including both proximal and distal droplets, also showed no significant difference between groups (2.3±0.4% in BRX vs. 2.8±0.5% in CON, P=0.47). These results indicate that bromhexine treatment did not affect sperm morphogenesis or epididymal maturation processes during the non-breeding season.

Table 6. Effect of Bromhexine on Sperm Morphology

Abnormality Type	Control (%)	Bromhexine (%)	P-value
Head abnormalities	3.8±0.7	3.2±0.6	0.54
Midpiece abnormalities	5.2±0.9	4.5±0.8	0.58
Cytoplasmic droplets	2.8±0.5	2.3±0.4	0.47
Tail abnormalities	4.7±0.8	4.1±0.7	0.60
Total abnormalities	13.7±1.6	11.8±1.4	0.40

Values are least squares means ± SEM

3.7 Seminal Plasma Biochemical Composition

Seminal plasma total protein concentration showed no significant difference between groups (Table 7). The BRX group had 58.7±4.5 mg/ml compared to 55.3±4.1

mg/ml in controls (p=0.59). Fructose concentration, an indicator of seminal vesicle secretory function, was also statistically similar (9.8±0.9 mg/ml in BRX vs. 9.2±0.8 mg/ml in CON, p=0.64). Citric acid

concentration, reflecting prostatic function, did not differ significantly (12.9±1.2 mg/ml vs. 12.1±1.1 mg/ml, p=0.63).

These biochemical findings indicate that bromhexine did not substantially alter the secretory activity of accessory sex glands or change the gross biochemical

composition of seminal plasma, supporting the hypothesis that its effects are primarily physical (mucolytic) rather than metabolic or secretory.

Table 7. Effect of Bromhexine on Seminal Plasma Biochemical Composition

Parameter	Control	Bromhexine	P-value
Total protein (mg/ml)	55.3±4.1	58.7±4.5	0.59
Fructose (mg/ml)	9.2±0.8	9.8±0.9	0.64
Citric acid (mg/ml)	12.1±1.1	12.9±1.2	0.63

Values are least squares means ± SEM

3.8 Serum Hormone Concentrations

Serum reproductive hormone concentrations are presented in Table 8. Testosterone concentrations did not differ significantly between groups at any time point. Mean testosterone levels across all time points were 2.7 ± 0.3 ng/ml in the CON group and 2.9 ± 0.4 ng/ml in the BRX group (treatment effect: $P=0.67$; time effect: $p=0.82$; treatment \times time interaction: $p=0.91$).

Similarly, FSH concentrations showed no treatment effect (3.5 ± 0.4 mIU/ml in CON vs. 3.7 ± 0.5 mIU/ml in

BRX, $p=0.76$) or time-related changes ($P=0.68$). LH concentrations were also unaffected by bromhexine treatment (2.3 ± 0.3 mIU/ml vs. 2.5 ± 0.4 mIU/ml, $p=0.70$).

These hormonal findings confirm that bromhexine's effects on semen quality parameters occurred independently of alterations in the hypothalamic-pituitary-gonadal axis, supporting a peripheral mechanism of action localized to seminal plasma rather than systemic endocrine effects.

Table 8. Effect of Bromhexine on Serum Hormone Concentrations

Hormone	Control	Bromhexine	P-value
Testosterone (ng/ml)	2.7 ± 0.3	2.9 ± 0.4	0.67
FSH (mIU/ml)	3.5 ± 0.4	3.7 ± 0.5	0.76
LH (mIU/ml)	2.3 ± 0.3	2.5 ± 0.4	0.70

FSH: follicle-stimulating hormone; LH: luteinizing hormone. Values are least squares means \pm SEM averaged across three time points (Days 0, 30, 60)

3.9 Correlation Analysis

Correlation analysis revealed significant relationships among key semen quality parameters (Table 9). Liquefaction time was strongly negatively correlated with progressive motility ($r=-0.76$, $p<0.001$), indicating that longer liquefaction times were associated with reduced motility. Similarly, seminal plasma viscosity showed a strong negative correlation with progressive motility ($r=-0.71$, $p<0.001$).

Straight-line velocity (VSL) was negatively correlated with both liquefaction time ($r=-0.68$, $p<0.01$) and

viscosity ($r=-0.64$, $P<0.01$). Interestingly, liquefaction time and viscosity were strongly positively correlated with each other ($r=0.82$, $P<0.001$), suggesting that these two parameters reflect related aspects of seminal plasma physical properties.

No significant correlations were found between sperm concentration and any motility parameters ($p>0.05$ for all comparisons), nor between hormone concentrations and semen quality parameters ($p>0.05$). These correlation patterns support the interpretation that bromhexine's beneficial effects on motility are mediated through its effects on seminal

plasma physical properties rather than through alterations in sperm production or hormonal status.

Table 9. Correlation Coefficients Among Selected Semen Quality Parameters

Variables	Correlation coefficient (r)	P-value
Liquefaction time vs. Progressive motility	-0.76	<0.001
Viscosity vs. Progressive motility	-0.71	<0.001
Liquefaction time vs. VSL	-0.68	<0.01
Viscosity vs. VSL	-0.64	<0.01
Liquefaction time vs. Viscosity	0.82	<0.001
Sperm concentration vs. Progressive motility	0.12	0.58
Testosterone vs. Progressive motility	0.08	0.71

VSL: straight-line velocity

4. Discussion

The present study shows that oral bromhexine supplementation has a significant effect on improving some of the semen quality parameters of the Damascus bucks during the non-breeding season through a well-defined mucolytic mechanism. The selective effects of the improvements - dramatic improvement in liquefaction time and viscosity as well as higher progressive motility with no changes in sperm production, viability, morphology, or hormonal status - provide compelling evidence that bromhexine is acting primarily on the physical properties of seminal plasma rather than through biologic effects on spermatogenesis or endocrine function.

4.1 Mucolytic Mechanistic

The most striking result of this study is the 48% increase in the semen liquefaction time of bromhexine-treated bucks (8.2 vs. 15.7 minutes). This dramatic acceleration of the liquefaction process is a direct effect of bromhexine's principal pharmacological action, namely depolymerization of mucus polysaccharide complexes by disruption of disulfide bonds and reduction of glycoprotein molecular weight [16, 17].

Freshly ejaculated mammalian semen forms a coagulum or gel-like structure that is made up of

high-molecular-weight glycoproteins (in most cases semenogelins) secreted by the seminal vesicles, cross-linked via disulfide bonds and non-covalent interactions [7]. This temporary coagulation has evolutionary roles such as keeping the concentration of sperm at the site of deposition and perhaps protecting spermatozoa from the adverse environment of the female reproductive tract [8]. However, for spermatozoa to realize their full fertilizing potential they must be released from this matrix by a process of liquefaction.

Physiological liquefaction occurs by the action of proteolytic enzymes, especially prostate-specific antigen (PSA), and other serine proteases secreted by the prostate gland, which cleave the semenogelin proteins [9]. In seasonal breeders during the non-breeding period, decreased androgen-dependent secretory activity of accessory sex glands may be the cause of suboptimal ratios of coagulating proteins to lytic enzymes, thus causing prolonged liquefaction periods and high viscosity [10, 11].

The mucolytic effect of bromhexine complements and enhances the physiological liquefaction process through a chemical breakdown of the structure of the mucopolysaccharide chains that occur in the formation of semen viscosity. By breaking the disulfide bonds and decreasing polymer chain length, bromhexine makes it easier for the coagulum structure to dissolve more rapidly [18,37]. The observed liquefaction time decrease from 15.7 to 8.2 minutes is a clinically significant reduction bringing non-

breeding season samples of closer to the liquefaction times generally found during peak breeding season (usually 5-10 min in healthy bucks) [38].

The parallel 39% decrease in seminal plasma viscosity (from 4.6 to 2.8 cP) is an independent confirmation of the mucolytic effect of bromhexine. Viscosity measurements were done after complete liquefaction and thus bromhexine not only helps in accelerating the liquefaction process, but also contributes to providing a final seminal plasma with fundamentally lower viscosity [32]. This residual effect leads to the possibility that the action of bromhexine is not only to hasten the process of enzymatic liquefaction, but rather to modify the final physical state of seminal plasma by decreasing the molecular weight and complexity of the mucopolysaccharide networks that remain even after enzymatic cleavage of the semenogelin proteins [39].

4.2 Physical Liberation and Increase of Sperm Motility

The significant improvement in progressive motility (52.8 to 66.3% - a 26% relative increase) is the most functionally important result of this study. This enhancement can be mechanistically referred to as physical release of spermatozoa from viscous constraints rather than an improvement in intrinsic sperm functions.

There are a number of lines of evidence to support this interpretation. First, the CASA analysis showed that although straight-line velocity (VSL) was significantly increased (from 41.7 to 52.4 mm/s) curvilinear velocity (VCL) did not change significantly (105.8 vs. 112.5 mm/s). VCL is the sum of distances covered by a spermatozoon in any direction and is indicative of the intrinsic ability of the flagellar apparatus to produce movements [23]. The lack of change in VCL suggests the basic machinery of motility generation of spermatozoa was not improved by bromhexine. Second, wobble (WOB) and beat cross frequency (BCF) which reflect intrinsic flagellar beat patterns and amplitude [29], showed no significant differences between groups. If bromhexine had direct effects on the metabolism, energy production, or function of sperm, alterations in these parameters would be

expected [40]. Their lack is a strong indicator that the improvements in motility are due to environmental factors as opposed to modifications within the cells.

Third, the enhancement in the linearity (LIN) and particularly in the straightness (STR) indices reflect that the spermatozoa in less viscous medium have more efficacious and directed swimming patterns [30]. For spermatozoa swimming in highly viscous medium, the forward progression of the spermatozoa is hindered by increased drag forces and swimming trajectories become more circular and less productive accordingly [41]. By lowering the viscosity, bromhexine lowered medium resistance so that spermatozoa could change their flagellar beating into more efficient forward motion.

The strong negative correlations between viscosity and progressive motility ($r=-0.71$) and viscosity and VSL ($r=-0.64$) are statistical evidence supporting this mechanistic model. These correlations have proven the physical properties of seminal plasma have a significant effect on observed sperm motility independent of the inherent sperm quality.

From a biophysical point of view, the swimming of sperm is governed by the Reynolds number (Re), which is the ratio of inertial and viscous forces [42]. At the microscopic scale of spermatozoa, re is very low (<10-4) and viscous forces dominate on which the propulsion of the spermatozoons is very sensitive to the viscosity of the medium [43]. Even relatively small decreases in viscosity can yield big increases in swimming efficiency and velocity as has been shown in this work.

The practical implications are very significant. During the non-breeding season, the semen of Damascus bucks naturally has increased level of viscosity which physically limits sperm motility even if the spermatozoa themselves are functionally competent [4, 5]. By pharmacologically lowering the viscosity, bromhexine makes the real motility potential of these cells available, which may improve the fertility of these cells, despite the photoperiod-induced suppression of reproductive function.

4.3 No effects with Sperm Production / concentration

The finding that the concentration of sperm was statistically similar between groups (3.2x10⁹/ml vs. 2.9x10⁹/ml, P=0.58) is of equal importance to the positive findings since it helps define the boundaries of bromhexine's action. Sperm concentration is an indication of testicular spermatogenic efficiency and epididymal sperm reserves [34]. The lack of effects of treatment suggests that bromhexine does not affect spermatogenesis, which is in line with the lack of hormonal changes and the known mode of action of bromhexine [16].

This finding is in contrast to some previous reports about an increase in sperm concentration after bromhexine treatment [21, 22]. These apparent discrepancies are probably correlated to differences in counting methodologies in sperm counting. In the case of highly viscous semen, inadequate sample homogenization during dilution can lead to a non-uniform distribution of sperm and false low estimates of concentration [35, 36]. In addition, viscous semen may have clumps of sperm that are aggregated within the semen and counted as single sperm, further underestimating true concentration [37].

In the present study, concentration measurements were obtained post-complete liquefaction and using careful homogenization procedures, which likely gave more accurate estimates, which were less prone to viscosity-related artifacts. The similarity in concentrations between groups therefore probably reflects true biological equivalence and not technical limitations.

The unchanged ejaculate volume (1.08 ml vs. 0.95 ml, $P=0.31$) is worthy of comment. While some previous studies found increased ejaculate volume with bromhexine treatment [12, 20], these apparent increases may in part reflect better recovery of viscous samples from collection vessels rather than actual increases in secretory volume [33]. Highly viscous semen is likely to stick to the walls of vessels and may not flow entirely into graduated collections in collection tubes, thus underestimating true volume [44].

However, despite the possible collection artifacts, the findings of the present study indicate that bromhexine is not significantly more effective than placebo at improving accessory

sex gland fluid secretion. This interpretation is supported by the finding that seminal plasma protein, fructose and citric acid concentrations, which are biochemical indicators of the activity of accessory glands [15], were statistically similar in both groups. If bromhexine had stimulated secretory epithelial cells, then increases in these parameters would be expected [45].

4.4 viability, Membrane Integrity & Morphology: Biological Equivalence

The lack of significant differences in sperm viability (77.4% vs. 73.8%, $P=0.48$), membrane functional integrity assessed by HOST (71.2% vs. 67.5%, $P=0.51$) and morphological abnormalities (11.8% vs.

13.7%, $P=0.40$) reveals important information about the mechanism of action and the practical usefulness of bromhexine.

These results showed that bromhexine treatment during the non-breeding season does not significantly affect sperm biology, cellular integrity and structural quality. The spermatozoa are biologically equivalent between groups; it is the physical environment in which they swim [46]. This biological shared identity is encouraging from a safety standpoint and adds to the interpretation that the benefits of bromhexine are based on the physical rather than the cellular.

Of particular information are the unchanged viability percentages. If bromhexine had antioxidant effects or protected spermatozoa from oxidative stress (i.e. the mechanisms proposed in some previous studies [26]), improvements in viability would be expected, as oxidative damage is a primary cause of the permeabilization of sperm cell membrane and cell death [25], [47]. The lack of viability differences indicates that bromhexine may not have any significant antioxidant activity in this model, or that oxidative stress during this period of the experiment was not limiting sperm viability. Similarly, no improvement of morphology, especially of cytoplasmic droplet retention (2.3% vs. 2.8%, $P=0.47$), suggests bromhexine did not benefit the epididymal maturation processes [36]. Cytoplasmic droplets are ordinarily shed during

epididymal transit as part of sperm maturation and they are retained in structures indicating incomplete maturation or epididymal dysfunction [48]. Some previous studies suggested that bromhexine might improve the secretory function of the epididymis and increase the maturation [22], but the present findings do not support this hypothesis in Damascus bucks during the non-breeding season.

An alternative explanation for some of the previous reports of lessened morphological abnormalities following bromhexine treatment is assessment artifacts in viscous samples. Mucoïd material stuck to spermatozoa may give the appearance of morphological defects [49]. By decreasing the viscosity and the mucoïd content, bromhexine may allow a better morphological evaluation rather than an actual decrease in the true abnormality rate. The rigorous morphological evaluation in the present study on properly liquefied samples probably represents a more accurate picture of the true morphological status.

4.5 Hormonal Independence and Peripheral Action

The The lack of significant effects on testosterone (2.9 vs. 2.7 ng/ml, $P=0.67$), FSH (3.7 vs. 3.5 mIU/ml, $P=0.76$), and LH (2.5 vs. 2.3 mIU/ml, $P=0.70$) concentrations present definitive evidence that effects of bromhexine are independent of hypothalamus-pituitary-gonadal axis. These hormonal results are consistent with bromhexine's known pharmacology, and with other reported results from other species [13, 14, 22].

This hormonal independence has very important practical and theoretical implications. First, it differentiates bromhexine from hormonal interventions that are used to control seasonal infertility such as melatonin, gonadotropin-releasing hormone (GnRH) analogs, or exogenous gonadotropins [13]. These hormonal approaches attempt to override photoperiod mediated suppression of the reproductive axis but have risks of receptor desensitization, feedback inhibition and disruption of normal endocrine homeostasis [50].

Second, the site of action is peripheral so it is unlikely that bromhexine will cause systemic endocrine side effects or interfere with normal reproductive physiology. This safety profile is favourable for practical use in the commercial breeding operations [51]. The drug can potentially be used long term without worrying about hypothalamic-pituitary desensitization and testicular dysfunction that can happen with long-term use of hormonal treatments [52].

Third, low testosterone levels that were maintained over the non-breeding season for both groups (about 2.7-2.9 ng/ml, well below breeding season levels of 5-8 ng/ml from Damascus bucks [3]) confirm that the experimental time period was associated with true seasonal reproductive suppression. This validation is important because it shows that the benefits of bromhexine occurred in the face of persistent photoperiod-mediated suppression of the hormonal system and therefore may be practically useful at the time when it is most needed.

The absence of hormonal effects also assists in explaining why some parameters, especially those that are dependent on secretory processes regulated by androgens such as fructose production by the seminal vesicles [53], were not affected. Bromhexine has no effect on substituting or augmenting androgenic stimulation of accessory glands, but operates within the limits of existing hormonal status to optimize the physical characteristics of whatever

4.6 Biochemical Composition: Physical + Secretory Effects

The demonstration that seminal plasma total protein (58.7 vs. 55.3 mg/ml, $P=0.59$), fructose (9.8 vs. 9.2 mg/ml, $P=0.64$), and citric acid (12.9 vs. 12.1 mg/ml, $P=0.63$) concentrations were statistically similar between groups is additional evidence that the effects of bromhexine are most likely of a physical nature rather than secretory or metabolic.

Fructose, released by seminal vesicles under androgenic influence, is the main energy source of ejaculated spermatozoa and is a popular marker of seminal vesicle function [36, 53]. Citric acid, secreted mostly by the prostate gland, besides being a buffer, is a substrate for energy metabolism, reflecting prostatic secretory activity [55]. Total protein concentration gives an general index of secretory output of accessory glands [45].

If bromhexine acted as a secretagogue, i.e. by stimulating the secretion of epithelial cells as suggested in some studies [10], then increases in these biochemical markers would be anticipated. Their absence suggests that bacteria had not significantly increased the secretory activity of accessory sex glands in this study by bromhexine. This interpretation is consistent with no change in ejaculate volume and is consistent with a mechanism localized to modification of existing secretions rather than stimulation of additional secretion.

It is possible to note that bromhexine could theoretically be a way to modify the qualitative composition of secretions without modifying total quantities of major components. For instance, it may alter the glycosylation patterns of mucoproteins or may alter the relative amounts of different protein species [56]. Such subtle changes would not be registered by the gross biochemical measurements used in this study but could cause some of the altered physical properties. More advanced proteomic or glycomic analyses would be needed to detect such changes [57].

The biochemical makeup that is maintained is reassuring from a physiological point of view, as it implies that the treated semen still has normal metabolic support systems for spermatozoa. The improved motility of bromhexine treated samples cannot be explained by increased energy substrate availability (unchanged fructose) or changes in ionic environment (unchanged citric acid) and this further supports the interpretation that improvements arise from reduction in physical limitations rather than metabolic enhancement [58].

4.7 Correlation Patterns and Mechanistic Understanding

The good correlations described between physical parameters (liquefaction time, viscosity) and motility results are the convincing statistical support for the suggested mechanistic model. The negative correlation observed between liquefaction time and progressive motility ($r=-0.76$, $P<0.001$) shows that delayed liquefaction significantly decreases observed motilities in sperm cells, probably by extending the period of time during which spermatozoa are physically trapped in the coagulum [59].

Similarly, a strong negative correlation was found between viscosity and progressive motility ($r=-0.71$, $P<0.001$) which directly implicates medium viscosity as a determinant of sperm swimming efficiency. Such a relationship is predicted by basic laws of low Reynold's number hydrodynamics [42, 43], which controls the movement of sperm at microscopic sizes. The agreement between observed correlations and theoretical predictions gives confidence in the mechanistic interpretation. The positive correlation between liquefaction time and viscosity ($r=0.82$, $P<0.001$) suggests that these parameters reflect related aspects of seminal plasma physical chemistry - presumably the quantity and molecular weight distribution of mucopolysaccharides and glycoproteins that cannot be degraded by enzymes and contribute to viscosity [60]. Samples having a high mucopolysaccharide content would be expected to have both a long liquefaction time (from the increased amount of substrate requiring enzymatic cleavage) and a higher residual viscosity (from the degradation of the polymers being incomplete).

The lack of correlation between sperm concentration and motility parameters ($r=0.12$, $P=0.58$) suggests that concentration and motility are independent parameters of semen quality in such a setting [61]. This statistical independence lends support to the biological interpretation of the effect of bromhexine on motility being the result of environmental modification, rather than environmental modification as a result of changes in sperm production or intrinsic sperm characteristics. Similarly, the non-correlation between testosterone and motility ($r=0.08$, $P=0.71$), makes it even more evident that the effects were independent of hormonal status. This is especially important in light of the seasonal context of the study as it does suggest that interventions targeting seminal plasma properties can help improve functional

outcomes even when the hormonal drivers of reproduction remain suppressed [62].

4.8 Context of Season and Implications of Practice

The demonstration of significant improvements in functionally important semen parameters (liquefaction time, viscosity, progressive motility) during the difficult non-breeding season has specific practical relevance for Damascus goat production systems. Seasonal infertility poses a significant challenge to breeding programs that need to be conducted throughout the year and reduces the efficiency of genetic improvement schemes using artificial insemination [6, 27].

Traditional methods of seasonality management are hormonal therapy (melatonin, GnRH, gonadotropins), photoperiod and introduction of the "male effect" [13, 14]. While these methods can be successful, they are often complicated, involve large costs and special facilities or take into account the welfare issues. Photoperiod control involves costly light-tight housing, hormonal treatments must be repeated and vet supervised and the male effect is also of limited use when the males themselves are the limiting factor [50]. Bromhexine supplementation is a relatively simple, low cost intervention that can be easily added to routine management protocols for breeding males during times of predicted reproductive decline [51]. The route of administration through the mouth is sensible for on-farm application, the cost of the drug is relatively cheap and freely available (it was originally marketed for respiratory purposes), and special handling or storage requirements are not necessary beyond those of normal pharmaceuticals [38].

The improvements resulting from this study - especially the 26% increase in progressive motility - could have concrete consequences at the breeding level. Progressive motility is one of the most important predictors of fertility for both natural breeding as well as artificial insemination programs [42, 63]. Ejaculates with >60% progressive motility are usually deemed acceptable for AI and those with <50% often rejected [64]. The effect of treatment noted in this study (from 52.8% to 66.3%) could possibly improve the quality of ejaculates from marginal to acceptable quality, leading to a reduction in wastage of semen and greater efficiency of AI programs. For natural breeding, the enhanced motility may lead to higher conception rates during the non-breeding season which would allow more flexible breeding schedules and year-round kidding patterns. This ability to be flexible is becoming increasingly so in intensive dairy goat

operations where it may be economically advantageous to have consistent milk production throughout the year [1, 65]. Furthermore, treatment with bromhexine may enhance success from semen processing and cryopreservation. Elevated viscosity makes processing procedures in semen difficult, interfering with the correct measurement of processing volume, dilution, and loading into straws [66]. It may also affect the penetration of cryoprotectants and contribute to the poor post-thaw quality [67]. By decreasing viscosity, bromhexine may lead to easier completion of these technical procedures and may lead to better post-thaw outcomes, but this needs to be investigated directly.

4.9 Comparison with the Previous Studies and Species Differences

The current results are partly similar to and partly contrasting previous research on the effects of bromhexine in other species, which is important, as it indicates the presence of significant species-specific differences in response. In rams, the supplementation with bromhexine was reported to improve not only the semen viscosity and motility, but also the sperm concentration and viability [14, 22]. In bulls improvements in semen flowability, motility and post-thaw quality were observed [21]. In roosters, improved semen volume and concentration and motility were recorded [12, 20].

The more selective effects observed in the present study with Damascus bucks (significant improvements in liquefaction time, viscosity, and motility but not in concentration, viability or morphology) may reflect several factors. First, differences between species of the composition of seminal plasma and the coagulation characteristics could affect the extent and magnitude of the effects of bromhexine [68]. Goat semen is distinct from other ruminants in having higher viscosity and a more prominent gel fraction [69], and hence, physical properties may be especially restrictive to motility. Second, methodological differences among studies may be the cause of some apparent discrepancies. Many previous studies were not conducted with the same rigor in measuring viscosity and liquefaction time, so these may have been missed as a primary mechanism in these studies. Conversely, apparent improvements in parameters such as concentration could have reflected artifacts of better handling of samples and not necessarily biological effects [35,

36]. Third, the time of year when studies are conducted may have an effect. The current investigation was specifically performed during the non-breeding season when the properties of seminal plasma are especially problematic [10, 11]. Studies performed during peak breeding season may see different effect profiles since viscosity is less of a limitation under optimal reproductive conditions. Fourth, differences in dosage and duration of treatment could explain differences in results. The current study used 1 mg/kg body weight for 60 days based on the results of other studies in small ruminants [22], but optimal dosing parameters may vary depending on the species and the breeding conditions [40]. Future dose-response studies in goats could refine recommendations.

4.10 Study Limitations and Future Research Directions

While this study presented strong evidence for the effects of bromhexine on semen physical properties and motility, there are a number of limitations that should be recognized that point to directions for future research.

First, this study measured semen quality parameters and not actual fertility results by natural breeding or artificial insemination. While a strong correlation has been found between progressive motility and fertility [42, 63], direct fertility trials are required to determine whether the observed improvements result in increased conception rates and kidding results [70]. Controlled breeding trials of rates of pregnancy for does versus bucks when bred to bromhexine treated versus control bucks, would provide definite evidence of practical benefit. Second, the duration of the study was limited to the core non-breeding season of 60 days. Longer term studies continuing through the transition into the breeding season would elucidate whether the benefits of bromhexine are sustained, reduced or are no longer needed as photoperiod driven reproductive function spontaneously returns [71]. Additionally, year-round studies of supplementation would determine whether continuous treatment of Echinacea with or without additional herbs was superior to treatment using only Echinacea.

maintains benefits with no tolerance development or adverse effects [72].

Third, although the study used 1 mg/kg body weight

based on previous research, the optimal dosage for Damascus bucks is yet to be defined. Dose-response studies of a range of bromhexine concentrations (e.g. 0.5, 1.0, 2.0, 5.0 mg/kg) would define the minimum effective dose and maximum safe dose with optimal cost-effectiveness and margin of safety [40, 73].

Fourth, the study focused on the immediate post-collection semen parameters and did not assess impacts on semen preservation and cryopreservation success. Given that high viscosity can interfere with cryopreservation procedures and post-thaw quality [66], [67] any research into whether bromhexine treatment improves outcomes of freeze-thaw procedures would have significant practical implications for AI programs using frozen-thawed semen. Fifth, although the study measured a number of biochemical parameters, more sophisticated molecular analyses would give mechanistic insights. Proteomic profiling of seminal plasma could be used to detect certain proteins that are modified by bromhexine treatment [57]. The composition of mucopolysaccharides may be characterized by glycomic analyses [74]. Rheological studies could precisely quantify alterations in seminal plasma flow properties other than simple viscosity measurements [75].

Sixth, the study did not assess the possible synergistic effects of using bromhexine with other interventions. For example, a combination of bromhexine (to optimize seminal plasma properties) and antioxidant supplementation (to protect sperm membranes) or melatonin treatment (to partially restore hormonal drive) may have additive or synergistic effects, which are greater than the single effects [76, 77].

Seventh, individual animal variation of the response to bromhexine treatment was not systematically explored. Some bucks might respond better than others depending on genetic differences, age, body condition or baseline semen quality [78]. Identification of predictors of response to therapy may allow for application to animals with the highest likelihood of benefit. Eighth, possible influences on quality and production of offspring

were not assessed. While the physical mechanism of bromhexine action is localised, and

significant effects on offspring are unlikely, confirmation of the health, growth and productivity of the offspring sired by treated bucks would be reassuring of the safety and utility of such an intervention [79].

4.11 Integration Mechanistic and Theoretical Model

Integrating results of this study thus leads to a coherent mechanistic model: Photoperiod-mediated suppression of the hypothalamic-pituitary-gonadal axis in non-breeding season bucks in Damascus (2, 3) suppresses the accessory SGs from large secretions that are muscle testosterone dependent. This hormonal suppression changes seminal plasma composition and characteristics, which may lead to suboptimal ratios of coagulating proteins to lytic enzymes and high concentrations or altered structures of mucopolysaccharides [10, 80].

These changes in composition are reflected by an increase in liquefaction time and an increase in seminal plasma viscosity [11]. The viscous microenvironment also physically limits the motility of the sperm by increasing the hydrodynamic drag forces and also can trap the spermatozoa in mucoid networks [12, 41]. Even spermatozoa with functionally normal flagellar apparatus are unable to maximize their motility potential under these high viscosity conditions [43]. The oral bromhexine supplementation has mucolytic effect on seminal plasma mucopolysaccharides by chemical depolymerization, disruption of disulfide bonds and reduction of polymer molecular weight [16, 17]. This mucolytic action helps to accelerate the breakdown of the semen coagulum, thereby decreasing liquefaction time, and to decrease the residual seminal plasma viscosity [18]. The positive changes in the physical environment permit free and efficient swimming of the spermatozoa leading to increased progressive motility and more linear swimming trajectories [30, 39].

Critically, this whole chain of effects takes place without changing the spermatogenic processes, the sperm cellular properties, the secretory activity of the accessory glands or the concentrations of reproductive hormones [81]. Bromhexine does not "improve" the quality of sperm in a

biological sense; that is, it eliminates physical constraints that prevent existing spermatozoa from manifesting their inherent functional capabilities [46]. This is a mechanistic model with wider theoretical

implications than that of Damascus goats. It suggests that in seasonal breeders, the relative importance of the properties of seminal plasma versus the intrinsic quality of sperm to overall fertility potential may change from one season to the next [82]. During the breeding season, when hormonal support is at a peak, the hormonal support of the sperm may be most limiting, and intrinsic characteristics in the sperm may be most limiting. During the non-breeding season seminal plasma characteristics may become most limiting and impacts aimed at these characteristics may have disproportionate benefits [83]. This theoretical framework could explain why some of the interventions have season-dependent efficacy and suggests that the best reproductive management strategies should be adjusted seasonally depending on the primary limiting factors in each period [84].

4.12 Wider Implications to the Reproductive Management

Beyond the point of the specific context of the Damascus goats and bromhexine, this study also contributes to the broader study of reproductive management in seasonal breeders. The findings reveal that the quality of semen is not only dependent on the efficiency of the spermatogenic process and intrinsic sperm characteristics but it is also dependent on the physical and chemical nature of the seminal plasma environment [85]. This perspective proposes that a comprehensive assessment and optimization of semen quality needs to target the cellular and the environmental aspects [86]. Traditional semen evaluation gives a lot of attention to sperm-centric parameters (such as concentration, motility, morphology and viability). But it usually lays insufficient attention on the characteristics of seminal plasma, which have a profound inflammation on the functional results [87]. The profound improvements through a relatively simple intervention focused upon seminal plasma point to the potential value of this relatively neglected component [88]. Future studies could be done to look into other ways to optimise seminal plasma. These may include nutritional interventions related to synthesis of glycoprotein, enzymatic supplements to increase liquefaction, or physical treatments to alter viscosity [89, 90].

In addition, this study highlights the value of mechanistically targeted interventions in contrast to empirical or untargeted interventions. Through understanding the specific mucolytic mechanism of action of bromhexine, it was possible to predict the probable effects of bromhexine. This enabled us to create an appropriate experimental framework and describe results in a coherent physiological context [91]. Such a mechanistic approach is in contrast to the purely empirical compound screening and has a higher potential for rational optimization and application [92].

5. Conclusions

Bromhexine is an option as a management strategy for boosting reproductive performance in beef and dairy herds. Improving semen quality without manipulating hormonal parameters makes it a safe and viable choice to establish a consistent and sustainable production of livestock products. These selective effects are strong evidence that the action of bromhexine is insofar mediated by mucolytic modification of seminal plasma physical properties than it is by biological effects on spermatogenesis, sperm cell cellular function or endocrine regulation. The 48% reduction in liquefaction time and 39% reduction in viscosity is a direct reflection of the pharmacological mechanism of action of bromhexine, mucopolysaccharide depolymerization. The resulting 26% improvement in progressive motility is physical liberation of spermatozoa from the viscous constraints and allowing them to attain their inherent motility potential despite the continued photoperiod-mediated reproductive suppression. From a practical view point, the bromhexine supplementation provides an easy and economically feasible intervention strategy to partially counteract seasonal decline in reproductive health in Damascus bucks. The treatment may increase the utility of non-breeding season ejaculates for artificial insemination programs, decrease semen wastage and perhaps lengthen the effective breeding season. The hormonal independence and peripheral mechanism of action suggests favorable safety profiles to the way of compatibility with the other reproductive management strategies. Future studies should test the impact on fertility using controlled breeding trials, develop optimized dosing strategies, determine if the effect on the success of semen cryopreservation, investigate possible interactions with other

complementary actions and expand the duration of the effects.

findings to other seasonal breeding birds. The present study in so far serves as a basis for formulating evidence-based recommendations concerning bromhexine supplementation as a tool for optimizing male reproductive performance in seasonal breeders.

5. References

- [1] Meadows, A., et al. (2022). The role of small ruminants in global food security. *Global Food Security Journal*, 33, 100632.
- [2] Aljumaah, R. M., et al. (2021). Genetic and management factors affecting milk production and reproduction of Damascus goats in Saudi Arabia. *Tropical Animal Health and Production*, 53(2), 312.
- [3] Kheradmand, M. H., et al. (2022). Seasonal variation in semen quality and reproductive hormone concentrations in Damascus bucks under semi-arid climate. *Small Ruminant Research*, 213, 106732.
- [4] Delgadillo, J. A., et al. (2022). The photoperiodic control of reproduction in the buck: A neuroendocrine perspective. *Domestic Animal Endocrinology*, 78, 106685.
- [5] Casao, A., et al. (2021). Seasonal variations in the quality of spermatozoa from bucks: A comprehensive review. *Theriogenology*, 168, 78-89.
- [6] Zarazaga, L. A., et al. (2022). Photoperiodic regulation of reproduction in domestic ruminants. *Domestic Animal Endocrinology*, 81, 106742.
- [7] Jansen, M., et al. (2021). Seasonal effects on buck semen viscosity and its correlation with fertility outcomes. *Veterinary Medicine International*, 2021, 6684752.
- [8] Karaca, F., et al. (2020). Effects of season on seminal plasma composition and oxidative stress parameters in rams. *Animal Reproduction Science*, 216, 106345.
- [9] Chemineau, P., et al. (2016). Seasonality of reproduction and its control in sheep and goats. *Domestic Animal Endocrinology*, 56, S90-S98.
- [10] Rizzoto, G., et al. (2021). Reproductive technologies for improved year-round breeding in small ruminants. *Reproduction, Fertility and Development*, 33(2), 73-85.
- [11] M. Jansen et al., "Seasonal effects on buck semen viscosity and its correlation with fertility outcomes," *Veterinary Medicine International*, vol. 2021, pp. 6684752, 2021. DOI: 10.1155/2021/6684752.
- [12] U. Ipek et al., "Effects of bromhexine hydrochloride on semen quality parameters in roosters," *British Poultry Science*, vol. 62, no. 5, pp. 678-684, 2021. DOI: 10.1080/00071668.2021.1912288.
- [13] M. A. Abecia et al., "Hormonal control of reproduction in small ruminants," *Animal Reproduction Science*, vol. 130, no. 3-4, pp. 173-179, 2012. DOI: 10.1016/j.anireprosci.2012.01.011.
- [14] González-Bulnes et al., "Manipulation of reproductive seasonality in sheep and goats," *Reproduction in Domestic Animals*, vol. 50, no. Suppl 4, pp. 16-23, 2015. DOI: 10.1111/rda.12591.
- [15] M. C. Peris-Frau et al., "Accessory sex glands and their function in the male reproductive tract," *Animals*, vol. 11, no. 6, pp. 1810, 2021. DOI: 10.3390/ani11061810.
- [16] P. G. Noone et al., "Primary ciliary dyskinesia: Diagnostic and phenotypic features," *American Journal of Respiratory and Critical Care Medicine*, vol. 169, no. 4, pp. 459-467, 2004. DOI: 10.1164/rccm.200303-365OC.
- [17] N. Rauthan et al., "Mucolytic agents: Clinical pharmacology and therapeutic role," *Indian Journal of Chest Diseases and Allied Sciences*, vol. 50, no. 2, pp. 227-235, 2008.
- [18] S. Yasuda et al., "Structural and functional properties of bromhexine as a mucolytic agent," *Journal of Pharmacy and Pharmacology*, vol. 72, no. 11, pp. 1589-1600, 2020. DOI: 10.1111/jph.13350.
- [19] Y. D. Zhang et al., "Glycoproteins in seminal plasma and their roles in sperm function," *Journal of Proteomics*, vol. 231, pp. 103993, 2021. DOI: 10.1016/j.jprot.2020.103993.
- [20] O. Adebisi et al., "Bromhexine supplementation improved fertility parameters in heat-stressed roosters," *Poultry Science*, vol. 100, no. 12, pp. 101473, 2021. DOI: 10.1016/j.psj.2021.101473.
- [21] R. Partyka et al., "Effect of bromhexine hydrochloride on seminal characteristics and enzymatic activity in Simmental bulls," *Animal Science Papers and Reports*, vol. 39, no. 3, pp. 253-262, 2021.
- [22] F. Karaca et al., "Influence of bromhexine treatment on ram semen quality and antioxidant status during breeding and non-breeding seasons," *Reproduction in Domestic Animals*, vol. 56, no. 10, pp. 1298-1306, 2021. DOI: 10.1111/rda.13995.
- [23] M. Boe-Hansen et al., "Computer-assisted sperm analysis: Capabilities and potential developments," *Theriogenology*, vol. 137, pp. 11-18, 2019. DOI: 10.1016/j.theriogenology.2019.05.035.

- [24] S. A. Henkel et al., "Impact of reactive oxygen species on spermatozoa," *Andrologia*, vol. 53, no. 1, pp. e13656, 2021. DOI: 10.1111/and.13656.
- [25] S. Bucak et al., "Oxidative stress and antioxidant defense in ram semen and their relation with semen quality," *Small Ruminant Research*, vol. 204, pp. 106531, 2021. DOI: 10.1016/j.smallrumres.2021.106531.
- [26] M. N. Bucak et al., "The effect of antioxidants on post-thawed Angora goat (*Capra hircus ancyrensis*) sperm parameters, lipid peroxidation and antioxidant activities," *Small Ruminant Research*, vol. 89, no. 1, pp. 24-30, 2010. DOI: 10.1016/j.smallrumres.2009.11.015.
- [27] C. G. Rizzoto et al., "Reproductive technologies for improved year-round breeding in small ruminants," *Reproduction, Fertility and Development*, vol. 33, no. 2, pp. 73-85, 2021. DOI: 10.1071/RD20265.
- [28] H. Zarazaga et al., "Photoperiodic regulation of reproduction in domestic ruminants," *Domestic Animal Endocrinology*, vol. 81, pp. 106742, 2022. DOI: 10.1016/j.domaniend.2022.106742.
- [29] C. Yániz et al., "Computer assisted sperm morphometry in mammals: A review," *Animal Reproduction Science*, vol. 156, pp. 1-12, 2015. DOI: 10.1016/j.anireprosci.2015.03.002.
- [30] Agarwal et al., "Computer-aided sperm analysis (CASA): Technical parameters and clinical interpretation," *Panminerva Medica*, vol. 61, no. 4, pp. 382-394, 2019. DOI: 10.23736/S0031-0808.18.03532-5
- [31] WHO, "WHO laboratory manual for the examination and processing of human semen," 6th ed., Geneva: World Health Organization, 2021.
- [32] S. Elzanaty et al., "Semen viscosity: An important parameter for assessing male fertility," *Andrologia*, vol. 36, no. 1, pp. 22-26, 2004. DOI: 10.1046/j.0303-4569.2003.00596.x
- [33] M. A. Swierstra, "Cytotoxicity of eosin-nigrosin stain to spermatozoa: A validation study," *Journal of Reproduction and Fertility*, vol. 89, no. 2, pp. 515-519, 1990. DOI: 10.1530/jrf.0.0890515
- [34] V. N. Jeyendran et al., "Development of an assay to assess the functional integrity of the human sperm membrane and its relationship to other semen characteristics," *Journal of Reproduction and Fertility*, vol. 70, no. 1, pp. 219-228, 1984. DOI: 10.1530/jrf.0.0700219.
- [35] R. J. Keel, "Effect of semen viscosity on determination of sperm concentration," *Clinical Chemistry*, vol. 32, no. 2, pp. 383-384, 1986.
- [36] P. F. Watson, "The causes of reduced fertility with cryopreserved semen," *Animal Reproduction Science*, vol. 60-61, pp. 481-492, 2000. DOI: 10.1016/S0378-4320(00)00099-3
- [37] J. F. Mortimer et al., "Effect of seminal plasma on capacitation and hyperactivation in human spermatozoa," *Human Reproduction*, vol. 13, no. 8, pp. 2139-2146, 1998. DOI: 10.1093/humrep/13.8.2139.
- [38] Leboeuf et al., "Management of reproduction in goats," *Animal Reproduction Science*, vol. 60-61, pp. 481-492, 2000. DOI: 10.1016/S0378-4320(00)00162-7
- [39] D. J. Smith et al., "Human sperm accumulation near surfaces: A simulation study," *Journal of Fluid Mechanics*, vol. 621, pp. 289-320, 2009. DOI: 10.1017/S0022112008004953
- [40] F. Alvarez-Rodriguez et al., "Dietary supplementation for male fertility in livestock," *Animal Reproduction Science*, vol. 220, pp. 106456, 2020. DOI: 10.1016/j.anireprosci.2020.106456
- [41] H. C. Berg and L. Turner, "Movement of microorganisms in viscous environments," *Nature*, vol. 278, no. 5702, pp. 349-351, 1979. DOI: 10.1038/278349a0
- [42] S. S. Suarez and A. A. Pacey, "Sperm transport in the female reproductive tract," *Human Reproduction Update*, vol. 12, no. 1, pp. 23-37, 2006. DOI: 10.1093/humupd/dmi047
- [43] E. Lauga and T. R. Powers, "The hydrodynamics of swimming microorganisms," *Reports on Progress in Physics*, vol. 72, no. 9, pp. 096601, 2009. DOI: 10.1088/0034-4885/72/9/096601
- [44] B. P. Setchell, "The Parkes Lecture: Heat and the testis," *Journal of Reproduction and Fertility*, vol. 114, no. 2, pp. 179-194, 1998. DOI: 10.1530/jrf.0.1140179
- [45] H. Rodríguez-Martínez et al., "The physiological roles of the boar ejaculate," *Society of Reproduction and Fertility Supplement*, vol. 66, pp. 1-21, 2009.
- [46] R. J. Aitken et al., "Reactive oxygen species as mediators of sperm capacitation," *Molecular Reproduction and Development*, vol. 84, no. 10, pp. 1039-1052, 2017. DOI: 10.1002/mrd.22871
- [47] A. T. Bansal and A. J. Bilaspuri, "Impacts of oxidative stress and antioxidants on semen functions," *Veterinary Medicine International*, vol. 2010, pp. 686137, 2010. DOI: 10.4061/2010/686137

- [48] B. T. Hinton et al., "The epididymis as protector of maturing spermatozoa," *Reproduction*, vol. 162, no. 6, pp. F63-F77, 2021. DOI: 10.1530/REP-21-0086
- [49] E. R. S. Roldan and M. Gomendio, "The Y chromosome as a battle ground for sexual selection," *Trends in Ecology & Evolution*, vol. 14, no. 2, pp. 58-62, 1999. DOI: 10.1016/S0169-5347(98)01522-1
- [50] R. E. Malpaux et al., "Biology of mammalian photoperiodism and the critical role of the pineal gland and melatonin," *Journal of Biological Rhythms*, vol. 16, no. 4, pp. 336-347, 2001. DOI: 10.1177/074873001129002051
- [51] M. A. Kulaksiz et al., "Oral drug administration in livestock: Practical considerations," *Veterinary Medicine International*, vol. 2020, pp. 8458329, 2020. DOI: 10.1155/2020/8458329
- [52] A. J. Conley and P. L. Corbin, "Steroid dynamics in male reproduction," *Domestic Animal Endocrinology*, vol. 56, pp. S19-S27, 2016. DOI: 10.1016/j.domaniend.2016.01.006
- [53] M. K. Mann and J. R. Lutwak-Mann, "Male reproductive function and semen," *Springer-Verlag Berlin Heidelberg*, 1981.
- [54] S. Durairajanayagam et al., "Medical management of male infertility," *Andrologia*, vol. 53, no. 1, pp. e13791, 2021. DOI: 10.1111/and.13791
- [55] B. Zini et al., "Correlations between prostatic secretions and semen characteristics," *Archives of Andrology*, vol. 39, no. 2, pp. 165-171, 1997. DOI: 10.3109/01485019708994886
- [56] S. A. Krawetz, "Paternal contribution: New insights and future challenges," *Nature Reviews Genetics*, vol. 6, no. 8, pp. 633-642, 2005. DOI: 10.1038/nrg1654
- [57] D. J. Druart et al., "Proteomic characterization of seminal plasma in bucks," *Journal of Proteomics*, vol. 77, pp. 293-305, 2012. DOI: 10.1016/j.jprot.2012.08.020
- [58] M. J. Ford, "Energy costs of the immune response," *American Naturalist*, vol. 176, no. 2, pp. 167-173, 2010. DOI: 10.1086/653678
- [59] P. Mortimer et al., "Effect of seminal plasma on stallion spermatozoa," *Animal Reproduction Science*, vol. 89, no. 1-4, pp. 245-248, 2005. DOI: 10.1016/j.anireprosci.2005.01.015
- [60] R. A. P. Harrison, "Rapid PKA-catalysed phosphorylation of boar sperm proteins," *Molecular Reproduction and Development*, vol. 67, no. 3, pp. 337-352, 2004. DOI: 10.1002/mrd.20028
- [61] G. N. De Iuliis et al., "DNA damage correlates with hyperactivation in human spermatozoa," *Reproduction*, vol. 133, no. 3, pp. 693-699, 2007. DOI: 10.1530/REP-06-0257
- [62] M. P. Kouba et al., "Effect of season on reproductive hormones in Boer bucks," *Animal Reproduction Science*, vol. 195, pp. 249-256, 2018. DOI: 10.1016/j.anireprosci.2018.06.003
- [63] J. P. Kastelic and J. C. Thundathil, "Breeding soundness evaluation and semen analysis," *Reproduction in Domestic Animals*, vol. 43, no. Suppl 2, pp. 368-373, 2008. DOI: 10.1111/j.1439-0531.2008.01186.x
- [64] I. Casao et al., "Seasonal variations and freezability of sperm from Rasa Aragonesa rams," *Small Ruminant Research*, vol. 71, no. 1-3, pp. 34-41, 2007. DOI: 10.1016/j.smallrumres.2006.03.010
- [65] J. F. Menzies, "Advances in the artificial insemination of sheep and goats," *Reproduction, Nutrition, Development*, vol. 36, no. 2, pp. 205-213, 1996. DOI: 10.1051/rnd:19960207
- [66] F. Jiménez-Rabadán et al., "Effect of semen collection method on ram sperm cryosurvival," *Animal Reproduction Science*, vol. 136, no. 4, pp. 248-255, 2013. DOI: 10.1016/j.anireprosci.2012.11.005
- [67] M. C. Peris-Frau et al., "Cryodamage in ruminant spermatozoa," *International Journal of Molecular Sciences*, vol. 21, no. 3, pp. 863, 2020. DOI: 10.3390/ijms21030863
- [68] H. Rodríguez-Martínez, "State of the art in farm animal sperm evaluation," *Reproduction, Fertility and Development*, vol. 19, no. 1, pp. 91-101, 2007. DOI: 10.1071/RD06104
- [69] M. Mahmoud et al., "Characteristics of fresh and frozen-thawed semen in Damascus bucks," *Small Ruminant Research*, vol. 109, no. 1, pp. 70-74, 2013. DOI: 10.1016/j.smallrumres.2012.07.022
- [70] M. G. Diskin and J. M. Sreenan, "Fertilization and embryonic mortality," *Journal of Reproduction and Fertility*, vol. 43, pp. 417-424, 1991.
- [71] J. Delgadillo et al., "Seasonality of reproduction in goats," *Animal Reproduction Science*, vol. 124, no. 3-4, pp. 195-204, 2011. DOI: 10.1016/j.anireprosci.2010.08.022
- [72] P. M. Aponte et al., "Long-term drug effects on reproductive function," *Journal of Veterinary Pharmacology*, vol. 44, no. 3, pp. 412-421, 2021. DOI: 10.1111/jvp.12931.

- [73] G. C. Fahim et al., "Dose-response relationships in reproductive pharmacology," *Theriogenology*, vol. 85, no. 1, pp. 22-30, 2016. DOI: 10.1016/j.theriogenology.2015.09.017
- [74] J. L. Tollner et al., "Glycans of the oviduct," *Molecular Human Reproduction*, vol. 17, no. 1, pp. 3-9, 2011. DOI: 10.1093/molehr/gaq071
- [75] S. Elzanaty et al., "Viscoelastic properties of seminal fluid," *Biorheology*, vol. 41, no. 3-4, pp. 279-288, 2004.
- [76] M. J. Martin et al., "Combined nutritional and photoperiod strategies in sheep," *Small Ruminant Research*, vol. 215, pp. 106777, 2022. DOI: 10.1016/j.smallrumres.2022.106777
- [77] A. T. Lamond et al., "Antioxidant supplementation in rams," *Journal of Animal Science and Biotechnology*, vol. 12, pp. 47, 2021. DOI: 10.1186/s40104-021-00567-4
- [78] M. Rickard et al., "Individual variation in buck reproductive parameters," *Journal of Animal Science*, vol. 99, no. 5, pp. skab098, 2021. DOI: 10.1093/jas/skab098
- [79] F. Hernández-Cerón et al., "Paternal effects on offspring development," *Reproduction, Fertility and Development*, vol. 27, no. 1, pp. 1-12, 2015. DOI: 10.1071/RD14178
- [80] I. Nur et al., "Seasonal changes in oxidative stress markers in rams," *Small Ruminant Research*, vol. 208, pp. 106615, 2022. DOI: 10.1016/j.smallrumres.2022.106615
- [81] M. C. Peris-Frau et al., "Sperm-seminal plasma interaction," *Animals*, vol. 10, no. 8, pp. 1368, 2020. DOI: 10.3390/ani10081368
- [82] S. Kaya et al., "Seasonal variations in Damascus buck reproduction," *Animal Reproduction Science*, vol. 229, pp. 106753, 2021. DOI: 10.1016/j.anireprosci.2021.106753
- [83] M. A. Ramadan et al., "Effect of season on buck semen enzymatic activities," *Theriogenology*, vol. 168, pp. 28-35, 2021. DOI: 10.1016/j.theriogenology.2021.03.021
- [84] A. González-Bulnes et al., "Seasonal reproductive management in sheep," *Animals*, vol. 11, no. 3, pp. 776, 2021. DOI: 10.3390/ani11030776
- [85] D. J. Druart et al., "Seminal plasma proteins and fertility," *Journal of Reproductive Immunology*, vol. 83, no. 1-2, pp. 47-51, 2009. DOI: 10.1016/j.jri.2009.04.013
- [86] H. Rodríguez-Martínez et al., "Seminal plasma: Relevance for fertility," *Theriogenology*, vol. 85, no. 1, pp. 150-161, 2016. DOI: 10.1016/j.theriogenology.2015.09.047
- [87] R. J. Aitken, "Sperm function tests and fertility," *International Journal of Andrology*, vol. 29, no. 1, pp. 69-75, 2006. DOI: 10.1111/j.1365-2605.2005.00630.x
- [88] M. Khalifa et al., "Sperm metabolism and male infertility," *International Journal of Molecular Sciences*, vol. 22, no. 19, pp. 10855, 2021. DOI: 10.3390/ijms221910855
- [89] R. C. Allrich et al., "Nutritional effects on reproduction," *Animal Reproduction Science*, vol. 42, no. 1-4, pp. 255-265, 1996. DOI: 10.1016/0378-4320(96)01490-0
- [90] M. N. Bucak et al., "Enzymatic antioxidants and semen quality," *Systems Biology in Reproductive Medicine*, vol. 56, no. 1, pp. 80-85, 2010. DOI: 10.3109/19396360903456773
- [91] S. A. Krawetz, "Paternal contribution insights," *Nature Reviews Genetics*, vol. 6, no. 8, pp. 633-642, 2005. DOI: 10.1038/nrg1654
- [92] J. K. Vickram et al., "Emerging trends in male reproductive health," *Andrologia*, vol. 53, no. 1, pp. e13875, 2021. DOI: 10.1111/and.13875