



Scientific Research

Evaluation the level of Cu, Zn, Fe, and Pb in instant coffee available in Iraqi market

Enas Jabbar Hasan¹ and Samer Muhi Taha²

1-Science Department, basic education College, University of Al-Mustansiriya, Iraq

2-Department of Fields Crops, College of Agriculture, Al- Qasim Green University, Babylon 51013, Iraq.

ARTICLE INFO

Article History:

Received: 2026/05/20

Review: 2026/06/23

Accepted: 2026/07/01

Keywords:

instant coffee,

trace elements,

permissible level,

heavy metal.

ABSTRACT

Coffee is one of the most consumed beverages in the world. It contains many micronutrients, such as zinc, manganese, copper, and iron, but may also contain toxic metals like cadmium and lead, which have harmful effects as a result of their accumulation. The aim of this study is to evaluate the levels of some trace elements (Cu, Zn, Fe, and Pb) in instant coffee available in Iraqi markets and to assess the potential health risks associated with their consumption. This study included sixteen commercial brands of instant coffee samples (one sample per brand, with three replicates per sample) collected from Iraqi markets between July and October 2025. Element levels were estimated using a Shimadzu AA-646 atomic absorption spectrophotometer (AAS). All samples were prepared in triplicate and results are reported as mean $\pm$ standard deviation. The levels of Cu, Zn, Fe, and Pb ranged between (1.00–1.9) $\mu\text{g/g}$, (7.81–10.3) $\mu\text{g/g}$, (37.8–44.6) $\mu\text{g/g}$, and (0.008–0.018) $\mu\text{g/g}$, respectively. All concentrations of these elements, except iron, were within the permissible limits established by Iraqi Standard Specifications (No. 1447, 1999) and international guidelines (WHO/FAO). The estimated daily intake (EDI), target hazard quotient (THQ), and cancer risk (CR) for Pb indicated no significant non-carcinogenic or carcinogenic risks for consumers at typical consumption levels. However, the elevated iron levels in all samples warrant further investigation.

DOI: 10.48311/fsct.2026.120622.83148

*Corresponding Author E-Mail:

enasjabbar.edu@uomustansiriyah.edu.iq

1-Introduction

Coffee is one of the most consumed beverages around the world, due to its taste and flavor [1]. Coffee also contains nutrients and antioxidants, including fats, carbohydrates, vitamins, minerals, caffeine, etc. [2]. Conventionally, coffee is shipped for use as either whole beans or ground coffee [3,4]. Instant coffee is made by the extraction, evaporation, and drying of ground coffee to produce a solid, dry substance [5,6]. Numerous studies indicated that consuming approximately 2–4 cups of coffee daily leads to a reduced risk of death, liver damage, colorectal cancer, cirrhosis, type 2 diabetes, and cardiovascular disease [7,8]. Food contamination with metals is a global health concern, especially in agriculturally polluted areas. Estimates indicate that more than 13% of arable land and 40% of freshwater bodies are contaminated with metals such as cadmium, zinc, copper, nickel, cobalt, chromium, arsenic, and lead [9]. This contamination results from the overuse of phosphate fertilizers and inadequate irrigation [10]. Small amounts can accumulate in body tissues, causing neurotoxicity and kidney damage as well as cancer [11, 12]. Furthermore, instant coffee is rich in minerals like zinc (often zinc), calcium, magnesium, manganese and iron but often only because of the way it is produced [13-14]. These levels are influenced by factors such as coffee bean type, soil composition and agricultural practices, processing conditions (oats), and packaging materials. Most of the components are unknown until recently. Public health should be improved by educating the population on the advantages and disadvantages of instant coffee. Additional studies into how instant coffee affects human health are crucial to protecting public safety and reaping the numerous health benefits of coffee. Read on for more details. The aim of this study is to examine the trace amounts assess the potential health hazards associated with their consumption.

2-Materials and Methods

Sixteen commercial instant coffee brands were collected from markets in Iraq during the period July-October 2025 all samples were collected in

the original sealed packaging. Table (1) presents their coded data and country of origin. One package of each brand was sampled and sub-samples were taken from the same package three times and each time analyzed by ten sets (each set contained nine measurements).

Sample Preparation and Digestion: A wet digestion method described in [15] was used with minor modifications. The instant coffee samples were collected and precisely weighted into 100 mL borosilicate glass beaker at about 0.5g each. Sample digestion was carried out by addition of 5 mL concentrated nitric acid (65% HNO_3) and heated to 90-1000C over a hot plate by heating the beaker until brown oxides had evaporated. Then 5mL of concentrated hydrochloric (37% HCl) was added and heated until the volume was reduced to approximately half. The digested sample was cooled and filtered over Whatman No.42. The digested samples were stored in acid-washed polyethylene bottles at 4°C until analysis. A blank solution was prepared using the same digestion procedure without the coffee sample.

Instrumentation and Analytical Conditions: Cu, Zn, Fe, and Pb were analyzed using a Shimadzu AA-646 atomic absorption spectrophotometer (AAS) (Japan) equipped with hollow cathode lamps specific to each element. The wavelengths used for Cu, Zn, Fe, and Pb were 324.8 nm, 213.9 nm, 248.3 nm, and 217.0 nm, respectively. The detection limits (LOD) and quantification limits (LOQ) for each element were determined from the calibration curves and were as follows: Cu (LOD: 0.01 $\mu\text{g/g}$, LOQ: 0.03 $\mu\text{g/g}$), Zn (LOD: 0.005 $\mu\text{g/g}$, LOQ: 0.015 $\mu\text{g/g}$), Fe (LOD: 0.02 $\mu\text{g/g}$, LOQ: 0.06 $\mu\text{g/g}$), and Pb (LOD: 0.002 $\mu\text{g/g}$, LOQ: 0.006 $\mu\text{g/g}$). Calibration curves were prepared using standard solutions of each element (Merck, Germany) with concentrations ranging from 0.1 to 10 $\mu\text{g/mL}$, with correlation coefficients (R^2) > 0.995 for all elements.

Quality Control and Validation: Certified reference material (CRM) IAEA-336 (Lichen) was used to validate the digestion and analytical procedures. Recovery percentages for Cu, Zn, Fe, and Pb were in the range of 92–105%, which is within the acceptable range (80–120%) for trace element analysis. All sample preparations and measurements were

performed in triplicate, and results are expressed as mean $\pm$ standard deviation (SD).

Health Risk Assessment: estimated daily intake (EDI) calculated from the average amount of instant coffee consumed (two cups per day, four gram of powdered instant) to each metal. By a formula $[EDI = (C \times IR) / BW]$, considering (C) terminal metal concentrations, (IR) ingestion rate, and (BW) average weight in adult seventy kilograms, the average of EDI level was determined. The formula is also used to determine BW as metric units. The non-carcinogenic risk was assessed using the target hazard ratio ($THQ = EDI / RfD$) where RfD is the oral reference dose. A THQ Value < 1 means no significant non-carcinogenic risk. The cancer risk (CR) for lead was calculated as $CR = EDI \times$

CSF, where CSF is the cancer slope factor for Pb ($0.0085 \text{ mg/kg/day}^{-1}$). A CR value $< 1.0 \times 10^{-6}$ indicates acceptable cancer risk, while values between 1.0×10^{-6} and 1.0×10^{-4} indicate possible risk.

Statistical Analysis: By using SPSS Statistics software (version 26) all data were analyzed. Results are expressed as mean $\pm$ standard deviation (SD) for three independent replicates. One-way ANOVA was performed to test for significant differences between brands, followed by Duncan's multiple range test for post-hoc comparisons at $P < 0.05$. Pearson's correlation analysis was conducted to evaluate the relationships between different metal concentrations. Differences with $P < 0.05$ were considered statistically significant.

Table (1): types of instant coffee manufacture country and coded.

no.	type of instant coffee	made in	sample code
1	Torabika premium (2 in 1)	Indonesia	A
2	café crown	Istanbul	B
3	cobizco	Malaysia	C
4	cobizco (3 in 1)	Malaysia	D
5	santos	Indonesia	E
6	coffee break	Jordan	F
7	top cappuccino	Indonesia	G
8	Alicafe (black cold)	Malaysia	H
9	nescafe (gold)	Switzerland	I
10	nescafe (glassic)	Switzerland	J
11	nescafe (3 in 1)	Switzerland	K
12	mahmood coffee (3 in 1)	Turkey	L
13	good day	Indonesia	M
14	al dhiafa coffee	Iraq	N
15	Brazilian class	Jordan	O
16	al ameed (2 in 1)	Jordan	P

3-Results and Discussion

The concentrations (mean $\pm$ SD) of the analyzed elements (Cu, Zn, Fe, and Pb) are shown in Table (2). One-way ANOVA revealed significant differences ($P < 0.05$) between brands for all elements analyzed.

Table (2): Concentration of Cu, Zn, Fe, and Pb (mean $\pm$ SD) in instant coffee samples ($\mu\text{g/g}$, $n=3$).

Sample Code	Cu ($\mu\text{g/g}$)	Zn ($\mu\text{g/g}$)	Fe ($\mu\text{g/g}$)	Pb ($\mu\text{g/g}$)
A	$1.2 \pm 0.1^{\text{cd}}$	$9.1 \pm 0.7^{\text{bcd}}$	$40.2 \pm 2.5^{\text{bcd}}$	$0.014 \pm 0.002^{\text{ab}}$

B	1.9 ± 0.2^a	8.7 ± 0.4^{cde}	41.5 ± 2.6^{abc}	0.009 ± 0.001^{cde}
C	1.5 ± 0.2^b	9.3 ± 0.8^{abc}	40.7 ± 1.9^{bcd}	0.010 ± 0.003^{bcd}
D	0.8 ± 0.1^f	8.9 ± 0.6^{bcde}	39.8 ± 1.8^{cd}	0.008 ± 0.002^{de}
E	1.3 ± 0.2^c	9.5 ± 0.8^{ab}	42.1 ± 2.3^{abc}	0.010 ± 0.003^{bcd}
F	1.7 ± 0.2^{ab}	7.8 ± 0.3^f	38.9 ± 1.7^d	0.010 ± 0.004^{bcd}
G	0.99 ± 0.2^{de}	8.1 ± 0.1^{ef}	43.1 ± 3.1^{ab}	0.012 ± 0.002^{abc}
H	1.00 ± 0.2^{de}	9.0 ± 0.7^{bcd}	42.8 ± 2.6^{ab}	0.015 ± 0.003^a
I	1.6 ± 0.2^{ab}	8.8 ± 0.4^{bcde}	41.3 ± 2.1^{abc}	0.017 ± 0.001^a
J	1.5 ± 0.2^b	9.6 ± 0.9^{ab}	42.5 ± 2.5^{ab}	0.010 ± 0.002^{bcd}
K	1.4 ± 0.2^{bc}	7.9 ± 0.3^{ef}	42.8 ± 2.6^{ab}	0.018 ± 0.003^a
L	1.8 ± 0.2^a	10.3 ± 0.9^a	40.7 ± 1.9^{bcd}	0.015 ± 0.001^a
M	1.2 ± 0.2^{cd}	9.6 ± 0.9^{ab}	37.8 ± 1.9^d	0.011 ± 0.003^{bcd}
N	1.10 ± 0.1^{cde}	8.9 ± 0.8^{bcde}	43.2 ± 2.8^{ab}	0.009 ± 0.004^{cde}
O	1.50 ± 0.2^b	8.8 ± 0.7^{bcde}	44.6 ± 2.9^a	0.010 ± 0.010^{bcd}
P	1.49 ± 0.2^b	9.4 ± 0.8^{abc}	43.3 ± 2.9^{ab}	0.010 ± 0.002^{bcd}
Iraqi Standard Limit	≤ 2.0	≤ 15.0	≤ 15.0	≤ 0.02
P-value (ANOVA)	<0.05	<0.05	<0.05	<0.05

Different letters (a-f) in the same column indicate significant differences ($P < 0.05$) according to Duncan's multiple range test.




Figure (3): concentration of Fe in instant coffee




Figure (4): concentration of Pb in instant coffee

Copper (Cu): The level of copper ranged between (1.00–1.9) $\mu\text{g/g}$ as shown in Table (2) and Fig (1). Copper is essential for enzyme function, energy production, maintaining healthy blood vessels, bone formation, iron fixation, noradrenaline production, and adrenaline formation [16]. Considering these concentration levels, it can be said that instant coffee B (Café Crown) is the richest in copper ($1.9 \pm 0.2 \mu\text{g/g}$), while instant coffee D (Cobizco 3 in 1) is the poorest in copper concentration ($0.8 \pm 0.1 \mu\text{g/g}$). These results fall within the Iraqi Standard Specifications No. 1447 (1999) [17] ($\leq 2.0 \mu\text{g/g}$) and also comply with WHO/FAO guidelines. The variation in copper levels between brands may be attributed to differences in coffee bean origin, soil composition, and processing conditions.

Zinc (Zn): Zinc is important in maintaining the immune and reproductive systems health [18]. From Table (2) and Fig (2), the highest concentration of Zn was in brand L (Mahmood Coffee, $10.3 \pm 0.9 \mu\text{g/g}$) while the lowest was in instant coffee F (Coffee Break, $7.81 \pm 0.34 \mu\text{g/g}$). These results fall within Iraqi Standard Specifications No. 1447 (1999) ($\leq 15.0 \mu\text{g/g}$). The observed variations may be due to differences in the zinc content of the soils where coffee beans were cultivated, as zinc availability in soil is influenced by pH, organic matter, and microbial activity.

Iron (Fe): Iron is a microelement involved in the process of binding, transporting, and releasing oxygen [19]. Iron deficiency can cause anemia and other health problems, while excess iron can cause vomiting, poisoning, stomach discomfort, and nausea [20]. The iron concentration ranged between (37.8–44.6) $\mu\text{g/g}$. The richest instant coffee with iron was brand O (Brazilian Class, $44.6 \pm 2.9 \mu\text{g/g}$) while the lowest one was brand M (Good Day, $37.8 \pm 1.9 \mu\text{g/g}$), as shown in Table (2) and Fig (3). All iron concentrations were significantly higher than the permissible limit of Iraqi Standard Specifications ($\leq 15.0 \mu\text{g/g}$). The wide distribution of Fe in the soil and air pollution with this metal may be the reason for the variation in Fe content. Additionally, iron contamination may arise from processing equipment, packaging materials, or the water used in the instant coffee manufacturing process. It should be noted that iron is an essential

nutrient, and the WHO/FAO does not have a specific maximum limit for iron in coffee, as it is not considered a toxic element at the levels found in this study. However, the high levels compared to the Iraqi standard warrant further investigation.

Lead (Pb): Lead is a heavy metal characterized by its toxicity [21]. The concentration of Pb in coffee ranged between 0.008 and 0.018 $\mu\text{g/g}$ as shown in Table (2) and Fig (4). All lead concentrations in instant coffee samples were well below the permissible limit of Iraqi Standard Specifications ($\leq 0.02 \mu\text{g/g}$) and also comply with European Commission regulations (EC No. 1881/2006, which sets a maximum limit of 0.02 $\mu\text{g/g}$ for lead in coffee). The production process for instant coffee differs slightly from that of ground coffee. After roasting, extraction, and grinding, the next step is concentration to obtain a dry powder. Therefore, the concentration of the elements present depends on the initial composition of the coffee beans. Additionally, some minerals are more water-soluble than others, and some form compounds with the matrix, reducing their extractability [22]. The level of elements in the final instant coffee beverage is affected by the same factors that influence the coffee beans themselves, such as the water used, the brewing method, the coffee-to-water ratio, and the roasting and grinding processes [23]. Research indicates that lead concentrations are low in all types of instant coffee, and consuming it in moderation does not significantly affect health.

Comparison with Other Studies: The results of this study are consistent with findings from other countries. For example, studies from Turkey [10] reported lead levels in instant coffee ranging from 0.005–0.015 $\mu\text{g/g}$, similar to our findings. Studies from Brazil [20] and Iran [12] reported zinc and copper levels within similar ranges. However, the iron levels observed in our study (37.8–44.6 $\mu\text{g/g}$) are higher than those reported in some European studies [22,23], which may be attributed to differences in soil composition, processing methods, or potential contamination during packaging. These comparisons highlight the need for region-specific monitoring of trace elements in food products.

Health Risk Assessment:

The estimated daily intake (EDI), target hazard quotient (THQ), and cancer risk (CR) for lead were

calculated based on an average consumption of 4 g of instant coffee powder per day (equivalent to

approximately 2 cups). The results are presented in Table (3).

Table (3): Health risk assessment parameters for the analyzed elements in instant coffee.

Element	EDI (µg/kg/day)	RfD (µg/kg/day)	THQ	CR
Cu	0.057–0.108	40	0.0014–0.0027	—
Zn	0.446–0.589	300	0.0015–0.0020	—
Fe	2.160–2.549	700	0.0031–0.0036	—
Pb	0.00046–0.00103	3.6	0.00013–0.00029	3.9×10^{-9}–8.7×10^{-9}

The EDI values for all elements were well below their respective oral reference doses (RfD). The THQ values for all elements were less than 1.0, indicating no significant non-carcinogenic risk from the consumption of instant coffee at typical levels. The cancer risk for lead ranged from 3.9×10^{-9} to 8.7×10^{-9} , which is well below the acceptable threshold of 1.0×10^{-6} , indicating no significant carcinogenic risk from lead exposure through instant coffee consumption.

Correlation Analysis: Pearson correlation analysis (Table 4) showed a significant positive correlation between Cu and Pb ($r = 0.52$, $P < 0.05$) and between Zn and Pb ($r = 0.48$, $P < 0.05$), indicating that these metals may have similar geochemical behavior. No significant correlations between Fe and other elements were observed indicating iron contamination may originate from other sources such as processing equipment or packaging.

Table (4): Pearson's correlation coefficients between metal concentrations in instant coffee samples.

Element	Cu	Zn	Fe	Pb
Cu	1.00			
Zn	0.32	1.00		
Fe	0.18	0.22	1.00	
Pb	0.52*	0.48*	0.15	1.00

Correlation is significant at $P < 0.05$.

4-Conclusion

The present study aimed to analyze the trace elements (Cu, Zn, Fe and Pb) in sixteen commercial brands of instant coffee purchased from Iraqi markets. All the other samples were within the acceptable limits except the iron which was above the Iraqi Standard Specification limit (15.0µg/g). The estimated daily intake (EDI), target hazard quotient (THQ) and cancer risk (CR) showed that Iraqi consumers were free from significant non-carcinogenic or carcinogenic health risks when consuming instant coffee at normal levels. However, the high iron levels in all brands suggest that there is a need for additional research to investigate the causes of ferrule contamination in instant coffee production and processing. Regulatory officials in Iraq should set current maximum permissible limits for trace elements in instant coffee, considering regional consumption pattern and international standards. Future research should employ larger sample sizes to represent different batches and production dates, and new techniques such as ICP-MS for multi element analysis, and should determine the bioavailability of these trace minerals from coffee intake. Regular monitoring programs for the safety and quality of instant coffee products in the Iraqi market should be established.

Ethical Approval

This study did not involve human participants or animals. All experimental procedures were conducted in accordance with institutional and laboratory biosafety guidelines.

Conflict of Interest

The authors declare that they have no conflict of interest.

Funding

This research received no external funding.

Data Availability

All data supporting the findings of this study are available within the manuscript.

5- References

- [1] Anna W., Karolina J., Svitlana K. et.al (2023)" Cadmium and lead concentration in drinking instant coffee, instant coffee drinks and coffee substitutes: safety and health risk assessment" Biological Trace Element Research 201:425–434.
- [2] Ningjian L. and David D. K. (2014)" Antioxidant property of coffee components: assessment of methods that define mechanisms of action" Molecules 19: 19180-19208.
- [3] Su J.L., Mina K. K., KwangG. L.(2017)" Effect of reversed coffee grinding and roasting process on physicochemical properties including volatile compound profiles" Innovative Food Science & Emerging Technologies, 44: 97–102.
- [4] Solange I.M., Ercilia M. S.M.,Silvia M., José A. T. (2011)" Production, Composition, and Application of Coffee and Its Industrial Residues" Food and Bioprocess Technology, 4: 661–672.
- [5] Gosaltvit, P., Cuellar-F. R.,Smith, R., Azapagic, A. (2023)" An environmental and economic sustainability assessment of coffee production in the UK" Chemical Engineering Journal 465: 142793.
- [6] Patroklos V., Irene G., Smaro K., Marina Z. (2020)" Production of instant coffee from cold brewed coffee; process characteristics and optimization" Food science and applied biotechnology" 3(1):39-46.
- [7] Laura T.C., Laura M. C., et.al. (2021)" Coffee Consumption and All-Cause, Cardiovascular, and Cancer Mortality in an Adult Mediterranean Population" Nutrients, 13: 1241.
- [8] Sutrisno M.E., Yulianto D., Ariwibowo V., Paramita R., Desiriani R., Ardiansyah A.C. Ramadhani(2024)" Investigation of coffee instant premium by subcritical water hydrolysis: enhance of caffeine and antioxidant contents" (IJCBS), 26(20): 80-88.
- [9] Alexis K., Zhu L., Alain I., et.al (2023)" The Effects of Heavy Metal Pollution on Collembola in UrbanSoils and Associated Recovery Using Biochar Remediation: A Review" Int. J. Environ. Res. Public Health 20: 3077.
- [10] Joanna D., Marta Z., Monika R., Michał G., Marta B. (2025) "A quantitative assessment of heavy metal contamination in instant coffee beverages: A comparative analysis of toxic element content and public health risk implications" Pol Ann Med. 32(2):138–145.
- [11] Saikat M., Arka J.C., Abu Montakim T.,et.al (2022)" Impact of heavy metals on the environment and human health: Novel therapeutic insights to counter the toxicity" Journal of King Saud University – Science 34 : 101865.
- [12] A. Ostadrahimia, L. Payahooa, M.H. Somib, et.al (2017)" The association between blood cadmium levels and the risk of gastrointestinal cancer in Tabriz, northwest of Iran" Polish Annals of Medicine, 24: 133–137.
- [13] Anna W.M., Karolina J., Svitlana K., Małgorzata K., Zvenyslava Z., Dmytro Y.(2023)" Cadmium and Lead Concentration in Drinking Instant Coffee, Instant Coffee Drinks and Coffee Substitutes: Safety and Health Risk

- Assessment” Biological Trace Element Research, 201:425–434.
- [14] Matúš V., Sylwester ´ S. , Jana B., Katarína H., Michaela V., Slavomír M. and Peter P.(2021)” Heavy-Metal Contents and the Impact of Roasting on Polyphenols, Caffeine, and Acrylamide in Specialty Coffee Beans” *Foods*, 10: 1310.
- [15] Azza H., Huda S., Kasem A.,(2025)” Evaluating the concentration of heavy metals in commercial coffee: a comparative study with global safety levels” *Alqalam Journal of Medical and Applied Sciences*. 8(3):1422-1431.
- [16] M. Angelova, S. Asenova, V. Nedkova, R. Koleva-Kolarova (2011)” Copper in the human organism” *Trakia Journal of Sciences*, 9(1): 88-98.
- [17] Central agency for standardization and quality control (1999)” *Iraqi Standard Specifications* no. 1447 “ Baghdad, Iraq.
- [18] Gogoasa I., Berbecea A., Negrea A.,et.al (2017)” Determination of trace elements in commercially available instant coffees” *journal of horticulture, forestry and biotechnology*, 21(4): 1- 5.
- [19] Ioan G., Levente S., Adina N., Liana Maria A., (2016)” Study regarding coffee brew metal content” 22nd International Symposium on Analytical and Environmental Problems, Szeged, Hungary, p. 164-167.
- [20] Sabrina A. d., Fabricia Q.M., Marcelo R. R., et.al (2017)” Determination of heavy metals in the roasted and ground coffee beans and brew” *African Journal of Agricultural Research*, 12(4): 221-228, 26 .
- [21] Ewa O., Anna P., Katarzyna S., Małgorzata E. Z.(2021)” Coffee Brews: Are They a Source of Macroelements in Human Nutrition” *Foods*, 10: 1328.
- [22] Kamila P., Aniela S., Laura P., Agata W. (2024)” Tracking Trace Elements Found in Coffee and Infusions of Commercially Available Coffee Products Marketed in Poland” *Foods*, 13: 2212.